

Hybrubins: Bipyrrrole Tetramic Acids Obtained by Crosstalk between a Truncated Undecylprodigiosin Pathway and Heterologous Tetramic Acid Biosynthetic Genes


Zhilong Zhao,[†] Ting Shi,[‡] Min Xu,[‡] Nelson L. Brock,[‡] Yi-Lei Zhao,[‡] Yemin Wang,[‡] Zixin Deng,[‡] Xiuhua Pang,^{*,†} and Meifeng Tao^{*,‡}

[†]State Key Laboratory of Microbial Technology, School of Life Sciences, Shandong University, Jinan 250100, China

[‡]State Key Laboratory of Microbial Metabolism, Joint International Research Laboratory of Metabolic & Developmental Sciences, and School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200030, China

Supporting Information

ABSTRACT: Heterologous expression of bacterial artificial chromosome (BAC) clones from the genomic library of *Streptomyces variabilis* Snt24 in *Streptomyces lividans* SBT5 which carried a truncated undecylprodigiosin biosynthetic gene cluster led to the identification of hybrubins A–C. The hybrubins represent a new carbon skeleton in which a tetramic acid moiety is fused to a 2,2'-dipyrrrole building block. Gene knockout experiments confirmed that hybrubins are derived from two convergent biosynthetic pathways including the remaining genomic *red* genes of *S. lividans* SBT5 as well as the BAC encoded *hbn* genes for the production of 5-ethylidenetetramic acid. A possible biosynthetic pathway was also proposed.



Prodiginines are a group of red-colored tripyrrrole natural products including undecylprodigiosin (**1**),¹ its cyclic derivative streptorubin B,² and prodigiosin (**2**).³ Despite their wide range of biological activities, including uses as antibacterial, -fungal, and -malarial agents,⁴ prodiginines are best known for their cytotoxic and immunosuppressive potentials that supposedly manifest themselves via different modes of action.⁵ Moreover, synthetic prodigiosin analogues like PNU-156804⁶ and Bcl2-inhibitor obatclax (GX15-070)⁷ are being evaluated in phase I/II clinical trials for potential applications in the treatment of different types of cancer.^{5b,8}

The biosynthesis of **1** in *Streptomyces coelicolor* relies on the key intermediates 4-methoxy-2,2'-bipyrrrole-5-carbaldehyde (MBC) and 2-undecylpyrrole (Scheme 1).⁹ Initially, L-proline (ring A) is activated by RedM and loaded onto the PCP RedO.¹⁰ Oxidation by RedW is followed by transfer of the resulting pyrrole-2-carboxyl moiety to RedX and condensation with RedN-bound malonate.^{10,11} Subsequently, L-serine is incorporated under cyclization to the second pyrrole ring (ring B) by the oxoamine synthase domain of RedN.¹¹ Oxidation by RedV (as proposed) and methylation by RedI complete the assembly of MBC.¹² The 2-undecylpyrrole building block (ring C) is obtained from acetyl-CoA, malonyl-CoA, and glycine by a PKS/NRPS machinery (RedK/L/P/Q/R).¹³ RedH catalyzes the condensation of MBC and 2-undecylpyrrole to yield **1**.¹⁴ In the last step, streptorubin B is formed by RedG-catalyzed carbocyclization.^{12b} Interestingly, production of **2** by the *pig* gene cluster of *Serratia marescens* follows a similar convergent biosynthetic

logic.^{12a} In both pathways, rings A and B originate from the common bipyrrrole intermediate MBC. However, this building block is then connected with 2-methyl-3-amylypyrrole by PigC for the biosynthesis of **2**.^{12a}


The remarkable bioactivities of the prodiginines have promoted a multidisciplinary search for new analogues of **1** and **2**. For example, a mutasynthetic approach utilized the MBC production defective mutants *S. coelicolor* $\Delta redM$ ¹⁴ and *S. marescens* *pigH::57* bp.¹⁵ Prodiginine production by these strains is dependent on supplementation of exogenous MBC. Subsequently, feeding of synthetic MBC analogues allowed for production of derivatives of **1** and **2** under replacement of ring A with a phenyl, indolyl, furyl, or thienyl group.^{14,15} Interestingly, the relatively broad substrate specificity of PigC even allowed for the incorporation of the non-native substrate 2,4-dimethylpyrrole as a substitute for ring C, which ultimately led to the enzyme catalyzed production of obatclax.¹⁵

Streptomyces lividans is a close relative of *S. coelicolor* that also produces the red pigments **1** and streptorubin B.¹⁶ Recently, the $\Delta redKL$ mutants *S. lividans* SBT5 and SBT18 have been described.¹⁷ SBT5 lacks the ability to produce 2-undecylpyrrole but still contains all genes for the biosynthesis of MBC, and SBT18 is an SBT5-derived strain containing an additional copy of the global regulator gene cassette *afsRS_{da}* (Figure S1). In this study, we complemented the $\Delta redKL$ mutant *S. lividans* SBT5 with a genomic bacterial artificial chromosome (BAC) library of

Received: December 20, 2015

Published: January 22, 2016

Scheme 1. Biosynthesis of (a) the Prodiginine Precursor MBC and (b) 1 in *S. coelicolor* and 2 in *S. marescens*



Streptomyces variabilis Snt24, a strain recently isolated from Shennongjia (Eastern Hubei, China) forest soil, to screen for exconjugants that could restore red pigmentation by production of hybrid secondary metabolites.¹⁸ This effort led to the identification of the hybrubins which constitute a new group of prodiginine analogues in which ring C is replaced by a tetramic acid moiety. These “non-natural” natural products are hybrids of the *red* pathway from *S. lividans* and a tetramic acid pathway from *S. variabilis* Snt24.

A genomic BAC library of *S. variabilis* Snt24 with 576 clones and an average insert size of 100 kb was constructed according to a standard protocol (cf. the SI for details).¹⁹ The complete BAC library was transferred into *S. lividans* SBT5, and the resulting exconjugants were grown for 7 d on several test media at 30 °C. Accumulation of a red pigment, indicative of prodiginine production, was observed for six exconjugants (BAC clones 2B5, 3E4, 3G4, 4H8, 5H5, and 6D1) on R3 medium. The six BAC clones were subsequently introduced into *S. lividans* SBT18, and all six BAC clones conferred red pigmentation to the exconjugants of *S. lividans* SBT18 as well (Figure 1A). Comparative metabolic profiling of *S. lividans* SBT18/BAC by HPLC revealed three distinct peaks (A–C) that were neither present in a vector control strain nor in *S. variabilis* Snt24 (Figure 1B).

The three peaks were purified from an *S. lividans* SBT18/6D1 fermentation culture and named hybrubin A–C. HR-ESI-MS analysis (Figure S2) revealed molecular formulas of C₁₆H₁₅N₃O₃ (hybrubins A and B) and C₁₄H₁₁N₃O₄ (hybrubin C). The ¹H NMR data of hybrubin A contained two sets of signals at a ratio of 2:1 (A₁: major component; A₂: minor component). Each set consisted of signals for 15 protons, including six aromatic/olefinic protons, two methyl groups, and three exchangeable protons. The ¹³C NMR data featured eight quaternary carbons, six methine groups, and two methyl groups for both isomers. Comparison of the NMR spectra with literature data for prodiginines, revealed a common dipyrrole moiety and the signals for this fragment were assigned accordingly (ring A and B). Furthermore, COSY, HSQC, and HMBC data allowed for the elucidation of the remaining fragment as 5-ethylidenetetramic acid. HMBC signals supported that a methine group bridged rings B and C (Figure 1C and Figures S3 and S4). The NMR data of hybrubin B showed two isomers (B₁ and B₂) at a ratio of 1:1 akin to the spectra of A₁ and A₂ (Figure 1C and Figures S3 and S5). Theoretically, there are four diastereomers for the proposed structure of hybrubins A and B that differ in the configuration of the C6'-

Figure 1. Discovery and characterization of hybrubins. (A) Red pigmentation of *S. lividans* SBT18/BAC colonies on R3 medium. (B) Comparative metabolic profiling of *S. lividans* SBT18/BAC exconjugants by HPLC (detection at 500 nm). The compound eluting at 15 min is not related to the hybrubins. The ratio of hybrubins A–C varied for different batches of fermentations. Hybrubin C could not be detected in SBT18/SH5. (C) Structures of hybrubins A–C.

C3'' and the C5''-C6'' double bonds. NOE correlations in the NOESY spectra were used to assign the (*Z*Δ5'',6'')-configuration for the major diastereomer A₁ (1''-H-7''-H) and the (*E*Δ5'',6'')-configuration for B₁ (1''-H-6''-H). Two sets of signals were found for hybrubin C at a ratio of 3.5:1 (C₁: major component; C₂: minor component) (Figure 1C and Figures S3 and S6). An additional carbonyl group and the lack of signals for an ethylidene group designated hybrubin C as the 5-oxo analogue of hyrubins A and B (Figure 1C). The occurrence of multiple sets of peaks in the NMR spectra of prodiginines^{5a} and tetramic acids²⁰ has been reported in the literature. If the interconversion between two isomers is slow enough, for example, because it requires rotation of a C–C bond,²⁰ a second set of peaks arises in the respective NMR spectra. Therefore, the second diastereomer (A₂/B₂/C₂) observed in the NMR spectra of each hybrubin possibly represents the respective *E/Z* isomer regarding the C6'–C3'' bond. The ¹H NMR and ¹³C NMR data of hyrubins A₁/A₂, B₁/B₂, and C₁/C₂ are summarized in Tables S2–S4 respectively.

Sequencing of the hybrubin producing BAC clone 6D1 gave a 115 kb DNA sequence. Judging from restriction analysis with *Bam*HI and *Pvu*II, the six pigmentation-conferring BAC clones shared a ca. 41 kb overlapping region encompassing 32 genes (Figure S7). To localize the hybrubin gene cluster, BAC 6D1 was truncated by restriction digests with *Asc*I (pHZLL1, 40 kb of the overlapping DNA region), *Sca*I (pHZLL2, 21 kb), *Eco*RI (pHZLL3, 13 kb), and *Bam*HI (pHZLL4, 5 kb) (Figure S8). All plasmids except for pHZLL4 were able to confer hybrubin production to *S. lividans* SBT18 (Figure S9). Therefore, the genes involved in the hybrubin biosynthesis had to be located within the 13 kb fragment of pHZLL3. pHZLL3 encodes nine open reading frames *hbnA-H* (Figure 2A, Table S5) including a




Figure 2. Analysis of the *hbn* gene cluster. (A) Organization of *hbnA*–*hbnH* on the hybrubin production conferring plasmid pHZLL3. (B) HPLC analysis of single gene deletion mutants.

PKS (*hbnA*, domain architecture: KS-ACP-TE) and an NRPS (*hbnB*, C-A-PCP) with a predicted adenylation domain specificity for L-threonine.²¹ Additionally, genes for two ketoreductases (*hbnC*, *hbnE*), an oxoamine synthase (*hbnD*), a short-chain dehydrogenase/reductase (*hbnF*), a dehydratase (*hbnN1*), a discrete acyl carrier protein (*hbnG*), and a putative hydrolase (*hbnH*) are found on pHZLL3.


Analogous to the prodiginine biosynthesis, hyrubins could be the product of a convergent biosynthetic pathway. The

bipyrrole building block MBC would likely be provided by the genome-encoded *red* gene cluster of the host strain, whereas the assembly of 5-ethylidenetetramic acid would rely on some of the BAC-encoded Hbn proteins. To test this hypothesis, pHZLL3 was transferred to the MBC production impaired strain *S. lividans* CXF28 ($\Delta redMN$, Figure S1). No hybrubin could be detected in a crude extract of the *S. lividans* CXF28/pHZLL3 exconjugant (Figure S9) which confirmed the crucial role of the *red* pathway for the hybrubin biosynthesis. To determine the genes for the biosynthesis of the tetramic acid building block, nine in-frame deletion mutations ($\Delta hbnA$ – $\Delta hbnH$) of pHZLL3 were constructed (Figure S10–S12) and transferred to *S. lividans* SBT18. Hybrubin production was completely abolished in the $\Delta hbnA$ and $\Delta hbnB$ mutants and significantly impaired in the $\Delta hbnH$ mutant (Figure 2), which designated these three genes responsible for the biosynthesis of 5-ethylidenetetramic acid.

Ring closure in the biosynthesis of tetramic acids proceeds in many cases by Dieckmann cyclization. Four groups of enzymes are known for this reaction, in particular (i) the redox-incompetent reductase* (R*) domain of fungal tetramate synthases,²² (ii) the TE domain of bacterial polycyclic tetramate macrolactam synthases,²³ (iii) two proteins PyrD3 and PyrD4 of the pyrroindomycin gene cluster,²⁴ and (iv) a recently discovered dedicated Dieckmann cyclase.²⁵ However, homologues of neither of the above-mentioned proteins are encoded in the 13 kb *hbn* gene cluster. Although, a TE domain is found within HbnA, phylogenetic analysis revealed that it is not located in the same clade as the Dieckmann cyclase TE domains (Figure S13). Instead, the closest neighbor of HbnA-TE is TycC-TE, which catalyzes the macrolactamization in the tyrocidine pathway of *Coralloccoccus coralloides*.²⁶

A hybrubin biosynthetic pathway was proposed combining the bioinformatic analysis with experimental data (Scheme 2):

Scheme 2. Proposed Hybrubin Biosynthetic Pathway



Activation of L-threonine by HbnB-A and loading onto HbnB-PCP is followed by HbnA-KS-catalyzed condensation to a malonate extender unit to give HbnA-ACP-tethered 5-hydroxy-4-amino-3-oxohexanoate. As HbnA is lacking an AT domain, the loading of malonyl-CoA to HbnA-ACP needs to be accomplished by an acyltransferase from the host acting in *trans*.²⁷ Subsequently, HbnA-TE catalyzes the lactamization under release of 5-(1-hydroxyethyl)tetramic acid. Dehydration by HbnH would finish the assembly of 5-ethylidenetetramic acid. The residual amounts of hyrubins found in the $\Delta hbnH$ mutant could be explained by other enzymes substituting HbnH for the dehydration in the mutant, although slow

nonenzymatic loss of water also remains a possibility. The dipyrrole building block MBC is synthesized by the genome-encoded *red* gene cluster (Scheme 1). RedH might condensate the two building blocks to yield hyrubins A and B, given the previously established promiscuity of the enzyme.¹⁴ Finally, hyrubin C arises by further oxidation of hyrubins A and B.

In summary, the hyrubins were identified by heterologous expression of BAC clones from an *S. variabilis* Snt24 genomic library in the 2-undecylpyrrole production-deficient MBC producer *S. lividans* SBT5. The convergent hyrubin biosynthetic pathway utilizes both the genome-encoded truncated *red* gene cluster of SBT5 as well as the BAC-encoded *hbn* gene cluster originating from *S. variabilis*. To our knowledge, hyrubins are the first examples of chimeras combining the 2,2'-dipyrrole and the tetramic acid moiety. Both compound classes, prodiginines and tetramic acids, are rich sources of bioactive natural products despite the lack of obvious cytotoxicity or antibacterial activity of the hyrubins in initial bioactivity tests. The identification of the *hbn* gene cluster paves the way for the production of a variety of new hyrubin analogues with diverse tetramic acid substituents for bioactivity tests. These could be obtained in a combinatoric approach by expression of other (partial) tetramic acid biosynthetic gene clusters in *S. lividans* SBT5.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b03609.

Experimental procedures and spectroscopic data for all new compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: tao_meifeng@sju.edu.cn.

*E-mail: pangxiuhua@sdu.edu.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Tobias Kieser for helpful discussions. This work was supported by the NSF (31170084) and the MOST (973:2012CB721000) of China, the Science and Technology Commission of Shanghai Municipality (15JC1400401), the State Key Laboratory of Microbial Technology Open Projects Fund (Project NO.M2015-08), and the Leopoldina Fellowship Program (German National Academy of Sciences Leopoldina, LPDS 2013-12 to N.L.B.).

■ REFERENCES

- (1) (a) Wasserman, H. H.; Rodgers, G. C., Jr.; Keith, D. D. *Chem. Commun.* **1966**, 2, 825–826. (b) Wasserman, H. H.; Rodgers, G. C.; Keith, D. D. *Tetrahedron* **1976**, 32, 1851–1854.
- (2) (a) Gerber, N. N.; McInnes, A. G.; Smith, D. G.; Walter, J. A.; Wright, J. L. C.; Vining, L. C. *Can. J. Chem.* **1978**, 56, 1155–1163. (b) Haynes, S. W.; Sydor, P. K.; Corre, C.; Song, L.; Challis, G. L. *J. Am. Chem. Soc.* **2011**, 133, 1793–1798.
- (3) (a) Wasserman, H. H.; McKeon, J. E.; Smith, L.; Forgione, P. J. *J. Am. Chem. Soc.* **1960**, 82, 506–507. (b) Rapoport, H.; Holden, K. G. *J. Am. Chem. Soc.* **1962**, 84, 635–642.

(4) Papireddy, K.; Smilkstein, M.; Kelly, J. X.; Shweta; Salem, S. M.; Alhamadsheh, M.; Haynes, S. W.; Challis, G. L.; Reynolds, K. A. *J. Med. Chem.* **2011**, 54, 5296–5306.

(5) (a) Fürstner, A. *Angew. Chem., Int. Ed.* **2003**, 42, 3582–3603. (b) Stankovic, N.; Senerovic, L.; Ilic-Tomic, T.; Vasiljevic, B.; Nikodinovic-Runic, J. *Appl. Microbiol. Biotechnol.* **2014**, 98, 3841–3858.

(6) Mortellaro, A.; Songia, S.; Gnocchi, P.; Ferrari, M.; Fornasiero, C.; D'Alessio, R.; Isetta, A.; Colotta, F.; Golay, J. *J. Immunol.* **1999**, 162, 7102–7109.

(7) Nguyen, N.; Marcellus, R. C.; Roulston, A.; Watson, M.; Serfass, L.; Madiraju, S. R. M.; Goulet, D.; Viallet, J.; Bélec, L.; Billot, X.; Acoca, S.; Purisima, E.; Wiegman, A.; Cluse, L.; Johnstone, R. W.; Beauparlant, P.; Shore, G. C. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, 104, 19512–19517.

(8) Williamson, N. R.; Fineran, P. C.; Gristwood, T.; Chawrai, S. R.; Leeper, F. J.; Salmond, G. P. C. *Future Microbiol.* **2007**, 2, 605–618.

(9) Cerdeño, A. M.; Bibb, M. J.; Challis, G. L. *Chem. Biol.* **2001**, 8, 817–829.

(10) Thomas, M. G.; Burkart, M. D.; Walsh, C. T. *Chem. Biol.* **2002**, 9, 171–184.

(11) Stanley, A. E.; Walton, L. J.; Zerikly, M. K.; Corre, C.; Challis, G. L. *Chem. Commun.* **2006**, 42, 3981–3983.

(12) (a) Williamson, N. R.; Simonsen, H. T.; Ahmed, R. A. A.; Goldet, G.; Slater, H.; Woodley, L.; Leeper, F. J.; Salmond, G. P. C. *Mol. Microbiol.* **2005**, 56, 971–989. (b) Sydor, P. K.; Barry, S. M.; Odulate, O. M.; Barona-Gomez, F.; Haynes, S. W.; Corre, C.; Song, L.; Challis, G. L. *Nat. Chem.* **2011**, 3, 388–392.

(13) Mo, S.; Sydor, P. K.; Corre, C.; Alhamadsheh, M. M.; Stanley, A. E.; Haynes, S. W.; Song, L.; Reynolds, K. A.; Challis, G. L. *Chem. Biol.* **2008**, 15, 137–148.

(14) Haynes, S. W.; Sydor, P. K.; Stanley, A. E.; Song, L.; Challis, G. L. *Chem. Commun.* **2008**, 44, 1865–1867.

(15) Chawrai, S. R.; Williamson, N. R.; Salmond, G. P. C.; Leeper, F. J. *Chem. Commun.* **2008**, 44, 1862–1864.

(16) (a) Horinouchi, S.; Beppu, T. *Agric. Biol. Chem.* **1984**, 48, 2131–2133. (b) Meschke, H.; Walter, S.; Schrepf, H. *Environ. Microbiol.* **2012**, 14, 940–952.

(17) (a) Bai, T.; Yu, Y.; Xu, Z.; Tao, M. *J. Huazhong Agric. Univ.* **2014**, 33, 1–6. (b) Chen, L.; Wang, Y.; Guo, H.; Xu, M.; Deng, Z.; Tao, M. *Appl. Environ. Microbiol.* **2012**, 78, 4526–4528.

(18) Hopwood, D. A.; Malpartida, F.; Kieser, H. M.; Ikeda, H.; Duncan, J.; Fujii, I.; Rudd, B. A. M.; Floss, H. G.; Omura, S. *Nature* **1985**, 314, 642–644.

(19) Luo, M.; Wing, R. A. In *Plant Functional Genomics*; Springer, 2003; pp 3–19.

(20) Schobert, R.; Schlenk, A. *Bioorg. Med. Chem.* **2008**, 16, 4203–4221.

(21) (a) Stachelhaus, T.; Mootz, H. D.; Marahiel, M. A. *Chem. Biol.* **1999**, 6, 493–505. (b) Challis, G. L.; Ravel, J.; Townsend, C. A. *Chem. Biol.* **2000**, 7, 211–224.

(22) (a) Sims, J. W.; Schmidt, E. W. *J. Am. Chem. Soc.* **2008**, 130, 11149–11155. (b) Liu, X.; Walsh, C. T. *Biochemistry* **2009**, 48, 8746–8757.

(23) (a) Blodgett, J. A. V.; Oh, D.-C.; Cao, S.; Currie, C. R.; Kolter, R.; Clardy, J. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, 107, 11692–11697. (b) Lou, L.; Chen, H.; Cerny, R. L.; Li, Y.; Shen, Y.; Du, L. *Biochemistry* **2012**, 51, 4–6.

(24) Wu, Q.; Wu, Z.; Qu, X.; Liu, W. *J. Am. Chem. Soc.* **2012**, 134, 17342–17345.

(25) Gui, C.; Li, Q.; Mo, X.; Qin, X.; Ma, J.; Ju, J. *Org. Lett.* **2015**, 17, 628–631.

(26) Trauger, J. W.; Kohli, R. M.; Mootz, H. D.; Marahiel, M. A.; Walsh, C. T. *Nature* **2000**, 407, 215–218.

(27) Cheng, Y.-Q.; Tang, G.-L.; Shen, B. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, 100, 3149–3154.