

Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Relationship between acid tolerance and cell membrane in *Bifidobacterium*, revealed by comparative analysis of acid-resistant derivatives and their parental strains grown in medium with and without Tween 80

Xu Yang · Xiaomin Hang · Min Zhang · Xianglong Liu · Hong Yang

Received: 11 November 2014 / Revised: 10 January 2015 / Accepted: 29 January 2015 / Published online: 17 February 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract The acid tolerance is particularly important for bifidobacteria to function as probiotics because they usually encounter acidic environments in food products and gastrointestinal tract passage. In this study, two acid-resistant derivatives *Bifidobacterium longum* JDY1017dpH and *Bifidobacterium breve* BB8dpH, which displayed a stable acid-resistant phenotype, were generated. The relationship between acid tolerance and cell membrane was investigated by comparing the two acid-resistant derivatives and their parental strains grown in medium with and without Tween 80. The fold increase in acid tolerance of the two acid-resistant derivatives relative to their parental strains was much higher when cells were grown in medium with Tween 80 (10^4 – 10^5 -fold) than without Tween 80 (181- and 245-fold). Moreover, when cells were grown in medium with Tween 80, the two acid-resistant derivatives exhibited more C18:1 and cycC19:0, higher mean fatty acid chain length, lower membrane fluidity, and higher expression of *efa* gene encoding cyclopropane fatty acid synthase than their parental strains. No significant differences in cell membrane were observed between the two acid-resistant derivatives and their parental strains when cells were grown in medium without Tween 80. The present study revealed that, when cells were grown in medium with Tween 80, the significant fold increase in acid tolerance of the two acid-resistant derivatives was mainly ascribed to the pronounced changes in

cell membrane compared with their parental strains. Results presented here could provide a basis for developing new strategies of cell membrane modification to enhance acid tolerance in bifidobacteria.

Keywords Bifidobacteria · Acid tolerance · Cell membrane · Tween 80

Introduction

Bifidobacteria 10

⁸ CFU/g) to function as probiotics (Aureli et al. 2011; Marteau and Rambaud 1993). However, various acidic environments (e.g., the low pH of fermented dairy products in which bifidobacteria are added as probiotics and the low pH of the stomach) commonly reduce the survival of bifidobacteria, resulting in less than the recommended sufficient numbers reaching the intestine. Consequently, acid tolerance is recognized as a desirable property of potential probiotic bifidobacteria (Parvez et al. 2006). However, the acid tolerance of most bifidobacteria is weak (Matsumoto et al. 2004; Ruiz et al. 2011), limiting the application of bifidobacteria in probiotic products.

X. Yang · M. Zhang · X. Liu · H. Yang (✉)
State Key Laboratory of Microbial Metabolism, and School of Life Science and Biotechnology, Shanghai Jiao Tong University, 800 Dongchuan Rd., Shanghai 200240, People's Republic of China
e-mail: hongyang@sjtu.edu.cn

X. Hang · M. Zhang
Institute of Bio-medicine, Shanghai Jiao Da Onlly Company Limited, Shanghai 200233, People's Republic of China

To improve the acid tolerance of bifidobacteria, it is necessary to understand acid tolerance mechanisms. To survive in acidic environments, several responses are employed by bifidobacteria, including maintenance of pH homeostasis by H^+ -ATPase, production of NH_3 , regulation of global signaling systems, and general stress response (Jin et al. 2012; Sánchez et al. 2006, 2007; Ventura et al. 2004a; Waddington et al. 2010). These studies mainly focused on the self-response of bifidobacteria to acid stress or adaptation. However, it is not clear why different species or strains of bifidobacteria have different acid tolerance levels. Since the cell membrane is regarded as the most important natural defense for cells, cell membrane with different properties may result in different levels of environmental tolerance, including acid tolerance. Despite the high level of genetic conservation in the cell membrane, the role of the cell membrane in acid tolerance differed even among closely related bacteria (Broadbent et al. 2010; Revilla-Guarinos et al. 2014; Wu et al. 2012). Accordingly, the relationship between the cell membrane and acid tolerance has not been established in bifidobacteria.

The aim of the present study was to explore the differences in acid tolerance between acid-resistant derivatives and their corresponding parental strains at the cell membrane level. In this study, we compared the acid tolerance, membrane fatty acid composition, and membrane fluidity between two acid-resistant derivatives and their parental strains. In addition, the expression of gene encoding key synthase for cyclopropane fatty acids (CFA) was also compared between two acid-resistant derivatives and their parental strains.

Materials and methods

Bacterial strains and growth conditions

Four strains of *Bifidobacterium* were used in this study. *B. longum* JDY1017 (deposited in the China General Microbiological Culture Collection Center (CGMCC) under accession number CGMCC No. 9882) and *B. breve* BB8 (CGMCC No. 9881) were isolated from the human intestine. *B. longum* JDY1017dpH (CGMCC No. 8371) and *B. breve* BB8dpH (CGMCC No. 8370) were acid-resistant derivatives of strains *B. longum* JDY1017 and *B. breve* BB8, respectively. The “dpH” in the name of the acid-resistant derivatives stands for “derivatives isolated in low pH conditions.” All strains were maintained by subculturing 1 % inocula in de Man Rogosa Sharpe (MRS) medium supplemented with 0.05 % (*w/v*) L-cysteine (MRSC) and incubating for 16 h at 37 °C under anaerobic conditions (AnaeroGen™, Oxoid Ltd, Basingstoke, UK). The cultures were stored at 4 °C between transfers and were subcultured three times prior to experimental use.

Isolation and molecular identification of acid-resistant derivatives

To isolate acid-resistant derivatives, *B. longum* JDY1017 and *B. breve* BB8 were cultured in MRSC medium anaerobically at 37 °C overnight. Cells of each culture were washed in phosphate-buffered saline with 0.05 % (*w/v*) L-cysteine (pH 7.4) and then transferred to fresh MRSC medium adjusted to low pH (pH 3.5 for *B. longum* JDY1017 and pH 3.2 for *B. breve* BB8) with 6 N HCl and incubated anaerobically at 37 °C overnight, after which aliquots were spread on MRSC agar (pH 6.5) and incubated at 37 °C for 3–4 days under anaerobic conditions to recover possible acid-resistant derivatives. After repeating the above processes 52 and 65 times respectively, two acid-resistant derivatives *B. longum* JDY1017dpH and *B. breve* BB8dpH were obtained.

For identification of strains, genomic DNA was extracted according to the method of Ausubel et al. (2002), which was slightly modified by adding a cell disruption step with a Fast prep instrument (Thermo Fisher Scientific, USA) prior to the extraction procedure. In order to obtain 16S rRNA partial gene (900-bp fragment length), universal oligonucleotides were used (5'-ATAATGCGGCCGCACGGGCGGTGTGTTC-3' and 5'-TAATAGCGGCCGCAGCMGCCGCGGTAA TWC-3') as previously described (Vaugien et al. 2002). The PCR products were purified with TaKaRa DNA fragment Purification Kit (TaKaRa Biotechnology, Dalian, China), and then sequenced on an ABI-Prism 3730 automated sequencer (PE Applied Biosystems, USA). Molecular identification of strains was accomplished by analyzing the 16S rRNA gene sequences using RDP classifier software (version 2.2) in the GenBank database (<http://www.ncbi.nlm.nih.gov/blast>).

Evaluation of acid tolerance of bifidobacterial strains

Stationary growth phase cells grown in MRSC medium with Tween 80 (1 g/L) and without Tween 80 at 37 °C for 16 h anaerobically were collected by centrifugation (7600×*g*, 10 min), respectively, washed twice with phosphate-buffered saline with 0.05 % (*w/v*) L-cysteine (pH 7.4), resuspended in phosphate-buffered saline with 0.05 % (*w/v*) L-cysteine (adjusted to pH 3.5 for *B. longum* and pH 3.2 for *B. breve* with 6 N HCl, respectively), and then incubated at 37 °C under anaerobic conditions. After the acid challenge for 4 h, serial dilutions of cells were spread on MRSC agar. Plates were incubated at 37 °C for 3–4 days under anaerobic conditions, after which colony-forming units (CFU) were enumerated. Survival was calculated by dividing the number of CFU per mL after acid challenge by the number of CFU per mL at the time 0. The fold increase in survival of the acid-resistant derivatives versus their parental strains was then calculated as the survival of acid-resistant derivatives divided by the

survival of their corresponding parental strains. The experiments were performed in triplicate.

Stability evaluation of acid-resistant phenotype of the two derivatives

The stability of acid-resistant phenotype of the two acid-resistant derivatives *B. longum* JDY1017dpH and *B. breve* BB8dpH was determined according to the following procedure. Briefly, cultures of each acid-resistant derivative were transferred daily for 20 consecutive days in fresh MRSC medium, with 1 % inocula at each transfer. The cultures after each transfer were grown at 37 °C for 16 h anaerobically, and then used to test their survival in acidic conditions (pH 3.5, 4 h for *B. longum* JDY1017dpH, and pH 3.2, 4 h for *B. breve* BB8dpH)

strain identifiers (were)-957929r72(w)0(e)°C0-29128(T)1.39fo000038(m)0(e)999904-32.307929r72(w)0(e)s1029779(p)0(e)a990003

Table 1 Primers of *cfa* gene used for real-time quantitative RT-PCR

Organism	Primer sequence (F/R: 5'–3')	Product size (bp)	Amplification efficiency
<i>B. longum</i>	ATGTCCAACGAGTTCTACG CAGCCGATGTCCAGTAAG	167	0.98
<i>B. breve</i>	GTATCCGCAGGAATACTCAG GGCATAGAACTCGTTAGACA	165	1.02

60 °C for 2.5 min. The *cfa* gene expression fold change of acid-resistant derivatives versus their parental strains was determined according to $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen 2001), using 16S rRNA as the reference gene as previously described (Gueimonde et al. 2004). Individual RT-PCR reactions were performed in triplicate.

Statistical analysis

Student's *t* test was employed to investigate statistical differences using SPSS version 19.0 (IBM, USA). Differences with *P* value less than 0.05 were considered statistically significant.

Results

Isolation and identification of acid-resistant derivatives

Two acid-resistant derivatives *B. longum* JDY1017dpH and *B. breve* BB8dpH were successfully isolated from *B. longum* JDY1017 and *B. breve* BB8 after repeating the isolation procedures many times, respectively. The partial 16S rRNA gene of the two acid-resistant derivatives was found to have 100 % identity with their corresponding parental strains (data not shown).

The standard MRSC medium, which included routinely Tween 80, was frequently used for cultivation of bifidobacteria in numerous studies (Collado and Sanz 2007; Jin et al. 2012; Sánchez et al. 2007; Waddington et al. 2010). We therefore investigated the acid tolerance of the two acid-resistant derivatives and their corresponding parental strains when cells were grown in standard MRSC medium containing Tween 80. As shown in Fig. 1, after exposed to acidic conditions (pH 3.5, 4 h for *B. longum*, and pH 3.2, 4 h for *B. breve*), the survival of *B. longum* JDY1017dpH and *B. breve* BB8dpH was approximately 10^4 – 10^5 -fold higher than that of their corresponding parental strains, respectively. In addition, the two acid-resistant derivatives had a stable acid-resistant phenotype by daily cultivation for 20 consecutive days (Fig. 2). During the comparison analysis of acid tolerance of the two acid-resistant derivatives and their corresponding parental strains, a surprising finding was that

when cells were grown in medium without Tween 80, the survival of *B. longum* JDY1017dpH and *B. breve* BB8dpH was only 181- and 245-fold higher than that of their corresponding parental strains, after exposed to acidic conditions (pH 3.5, 4 h for *B. longum*, and pH 3.2, 4 h for *B. breve*), respectively (Fig. 1).

These results indicated that the fold increase in acid tolerance of the acid-resistant derivatives relative to their corresponding parental strains was more significant




Fig. 1 Fold increase of survival of the acid-resistant derivatives versus their corresponding parental strains grown in medium with and without Tween 80, respectively. **a** Fold increase of survival of *B. longum* JDY1017dpH versus its parental strain JDY1017 after exposed to pH 3.5 for 4 h. **b** Fold increase of survival of *B. breve* BB8dpH versus its parental strain BB8 after exposed to pH 3.2 for 4 h. Vertical lines on the bars represent standard deviations




Fig. 2 Stability of acid-resistant phenotype of the two acid-resistant derivatives. Cultures of each acid-resistant derivative and its parental strain were transferred daily for 20 consecutive days in fresh MRSC medium, respectively. The daily cultures were used to evaluate the survival in acidic conditions (pH 3.5, 4 h for *B. longum* JDY1017dpH and JDY1017, and pH 3.2, 4 h for *B. breve* BB8dpH and BB8), respectively. **a** Survival of *B. longum* JDY1017 and *B. longum* JDY1017dpH cells after exposure to pH 3.5 for 4 h. **b** Survival of *B. breve* BB8 and *B. breve* BB8dpH cells after exposure to pH 3.2 for 4 h. The error bars represent the standard deviations calculated from three independent experiments

in growth medium with Tween 80 than without Tween 80. Previous studies reported that Tween 80 was a derivative of octadecenoic acid, which as the exogenous fatty acid could be incorporated into cell membrane directly, and thus was closely related to cell membrane fatty acid composition in some lactic acid bacteria (Broadbent et al. 2014; Johnsson et al. 1995). Moreover, in *B. animalis*, different strains showed the differences in ability to incorporation of octadecenoic acid from the growth medium with Tween 80 into cell membrane (Oberge et al. 2013). These encouraged us to further compare cell membrane of the acid-resistant derivatives and their parental strains grown in MRSC medium with and without Tween 80.

Membrane fatty acid composition of the two acid-resistant derivatives and their parental strains grown in medium with and without Tween 80

The membrane fatty acid composition of the two acid-resistant derivatives and their parental strains grown in medium with and without Tween 80 was determined (Fig. 3 and Table 2), respectively. Seven main fatty acids were observed among all strains, dodecanoic acid (C12:0), tetradecanoic (C14:0), hexadecanoic acid (C16:0), hexadecenoic acid (C16:1), octadecanoic acid (18:0), octadecenoic acid (C18:1), and CFA (cycC19:0).

When cells were grown in medium with Tween 80, the membrane fatty acid composition of the two acid-resistant derivatives differed significantly from that of their corresponding parental strains (Fig. 3 and Table 2). Specifically, the acid-resistant derivatives displayed more C18:1 and




Fig. 3 Membrane fatty acid composition of the two acid-resistant derivatives and their corresponding parental strains grown in medium with and without Tween 80. **a** *B. longum* JDY1017 and *B. longum* JDY1017dpH. **b** *B. breve* BB8 and *B. breve* BB8dpH. Error bars correspond to the standard deviations of the mean, and different letters within each lipid type indicate that the means differ significantly ($P < 0.05$)

Table 2 Mean fatty acid chain length and ratio of saturated fatty acids to unsaturated fatty acids of the two acid-resistant derivatives and their parental strains grown in medium with and without Tween 80

	Medium with Tween 80				Medium without Tween 80			
	<i>B. longum</i> JDY1017	<i>B. longum</i> JDY1017dpH	<i>B. breve</i> BB8	<i>B. breve</i> BB8dpH	<i>B. longum</i> JDY1017	<i>B. longum</i> JDY1017dpH	<i>B. breve</i> BB8	<i>B. breve</i> BB8dpH
SFA/UFA	1.18±0.07	0.65±0.04*	0.62±0.03	0.28±0.01*	1.55±0.06	1.54±0.06	0.80±0.04	0.79±0.03
Mean fatty acid chain length	16.50±0.01	17.04±0.02*	16.92±0.02	17.48±0.01*	16.26±0.03	16.28±0.05	16.71±0.03	16.70±0.05

SFA total saturated fatty acids, UFA total unsaturated fatty acids, SFA/UFA the ratio of saturated fatty acids to unsaturated fatty acids without considering cyclopropane fatty acid (cycC19:0)

All values are means ± standard deviations. The asterisk represents significant difference between the two acid-resistant derivatives and their corresponding parental strains ($P<0.05$)

cycC19:0 content, less C14:0 and C16:0 content than their corresponding parental strains. These differences resulted in the lower SFA/UFA ratio and higher mean fatty acid chain length in the acid-resistant derivatives than those in their parental strains. The mean fatty acid chain length was 16.50 for *B. longum* JDY1017 and 17.04 for its acid-resistant derivative JDY1017dpH, and 16.92 for *B. breve* BB8, and 17.48 for its acid-resistant derivative BB8dpH. On the other hand, when cells were grown in medium without Tween 80, no significant differences in membrane fatty acid composition between the two acid-resistant derivatives and their corresponding parental strains were observed (Fig. 3 and Table 2). These results indicated that the addition of Tween 80 to the growth medium resulted in the significant differences in membrane fatty acid composition between the two acid-resistant derivatives and their corresponding parental strains.

Membrane fluidity of the two acid-resistant derivatives and their parental strains grown in medium with and without Tween 80

The membrane fluidity of the two acid-resistant derivatives and their parental strains grown in medium with and without Tween 80 was investigated by measuring the rotational diffusion of the fatty acyl chains. The steady-state fluorescence anisotropy values in stationary growth phase cells, which were estimated using DPH as a probe, are shown in Fig. 4. When cells were grown in medium with Tween 80, the two acid-resistant derivatives displayed higher anisotropy values than their corresponding parental strains ($P<0.05$), suggesting a lower membrane fluidity in the acid-resistant derivatives. Specifically, the anisotropy value was 0.126 for *B. longum* JDY1017, and 0.135 for its acid-resistant derivative JDY1017dpH, and 0.130 for *B. breve* BB8, and 0.138 for its acid-resistant derivative BB8dpH. However, when cells were grown in medium without Tween 80, no significant

differences in anisotropy values were observed between the two acid-resistant derivatives and their parental strains.




Fig. 4 Anisotropy values of the two acid-resistant derivatives and their corresponding parental strains grown in medium with and without Tween 80. **a** *B. longum* JDY1017 (white bars) and *B. longum* JDY1017dpH (gray bars). **b** *B. breve* BB8 (white bars) and *B. breve* BB8dpH (gray bars). Higher anisotropy values indicate lower membrane fluidity. The bars followed by asterisk indicate significant difference between the acid-resistant derivatives and their corresponding parental strains ($P<0.05$). Vertical lines on the bars represent standard deviations

Transcriptional expression of *cfa* gene

Previous studies reported that CFA synthase encoded by *cfa* gene could catalyze the conversion of C18:1 to CFA, and CFA was generally regarded as a tool to reduce membrane fluidity in many bacteria (Grogan and Cronan 1997; To et al. 2011). We therefore investigated the *cfa* gene expression fold change of the two acid-resistant derivatives relative to their corresponding parental strains by RT-PCR according to $2^{-\Delta\Delta Ct}$ method (Fig. 5). When cells were grown in medium with Tween 80, the expression of *cfa* gene was 3.6-fold higher in *B. longum* JDY1017dpH than that in *B. longum* JDY1017 and was 2.2-fold higher in *B. breve* BB8dpH than that in *B. breve* BB8. When cells were grown in medium without Tween 80, the expression of *cfa* gene displayed almost the same level between the acid-resistant derivatives and their corresponding parental strains.

Discussion

As part of strategies to improve acid tolerance in bifidobacteria, it is necessary to understand the mechanisms involved in development of such tolerance. Molecular genetic tools available for analysis of bifidobacteria are rather limited, strongly restricting investigations of their acid tolerance mechanisms (Berger et al. 2010; Sánchez et al. 2006, 2007; Ventura et al. 2004b). Therefore, the present study compared the cell membrane of acid-resistant derivatives and their parental strains in order to reveal the relationship of cell membrane with acid tolerance in bifidobacteria.

Strategies based on short exposures to sublethal acidic conditions were applied to improve acid tolerance of bifidobacteria, but strains with stable acid-resistant phenotype




Fig. 5 The *cfa* gene expression fold change of the two acid-resistant derivatives relative to their corresponding parental strains by RT-PCR according to $2^{-\Delta\Delta Ct}$ method. Cells were grown in medium with and without Tween 80, respectively. Vertical lines on the bars represent the standard deviations

were not always generated (Maus and Ingham 2003; Saarela et al. 2004). As an alternative strategy, wild-type strains were prolonged exposed to lethal acidic conditions for obtaining strains with stable acid-resistant phenotype (Collado and Sanz 2007). In the present study, two bifidobacterial acid-resistant derivatives were obtained by the method of prolonged exposure to lethal acidic conditions, exhibiting higher stable acid tolerance than their corresponding parental strains (Fig. 2). Additionally, the fold increase in acid tolerance of the acid-resistant derivatives relative to their corresponding parental strains when cells were grown in medium with Tween 80 (about 10^4 – 10^5 -fold) was much higher than that when cells were grown in medium without Tween 80 (181- and 245-fold) (Fig. 1). This suggested that the acid-resistant derivatives had a trait that utilizing Tween 80 in medium to exhibit greatly increase in acid tolerance.

In the present study, the two acid-resistant derivatives exhibited more C18:1 content than their corresponding parental strains when cells were grown in medium with Tween 80 (Fig. 3). Previous studies reported that some lactic acid bacteria such as *Lactobacillus* and *Lactococcus* could incorporate the exogenous C18:1 from growth medium containing Tween 80 (a derivative of C18:1) into cell membrane directly, consequently changing their fatty acid composition (Broadbent et al. 2014; Johnsson et al. 1995). Oberg et al. (2013) reported that the growth medium with Tween 80 resulted in an increase in C18:1 content of *B. animalis*. In *Lactobacillus casei*, more C18:1 content was observed in an acid-resistant derivative compared with the wild type (Wu et al. 2012). Furthermore, Fozo et al. (2004) reported that the mutant of *Streptococcus mutans* defective in production of monounsaturated fatty acids including C18:1 was extremely sensitive to acidic environments compared with the wild type, and the acid-sensitive phenotype of the mutant could be relieved by growth in the presence of exogenous C18:1. Taken together, the present results suggested that the two acid-resistant derivatives had a stronger ability to incorporate Tween 80 from medium than their corresponding parental strains, which consequently led to much more C18:1 content in the acid-resistant derivatives than in the corresponding parental strains when cells were grown in medium with Tween 80.

The C18:1 could be converted to CFA (cycC19:0) by the transfer of a methylene group from *S*-adenosyl-L-methionine to a double of unsaturated fatty acid chains of membrane phospholipids. This conversion was catalyzed by CFA synthase in many bacteria (Grogan and Cronan 1997; To et al. 2011). In the present study, the two acid-resistant derivatives displayed more cycC19:0 content and higher expression of *cfa* gene encoding CFA synthase than their corresponding parental strains when cells were grown in medium with Tween 80 (Figs. 3 and 5). More cycC19:0 content in the acid-resistant

derivatives could be ascribed to their more C18:1 content and higher expression of *cfa* gene. Additionally, previous studies reported that *cfa* defective mutants of *E. coli* and *Salmonella enterica* serovar *typhimurium* were more acid-sensitive (Chang and Cronan 1999; Kim et al. 2005). The CFA was generally regarded as a tool to prevent the entry of undesirable molecules (e.g., H⁺) into cells (Mykytczuk et al. 2007; Wu et al. 2012). It was also reported that the increased CFA content contributed to the enhancement of acid tolerance in *E. coli* (Brown et al. 1997), *Salmonella enteritidis* (Yang et al. 2014b), and *Lactobacillus casei* (Wu et al. 2012). Taken together, more CFA content could be one of the reasons for the higher acid tolerance of two acid-resistant derivatives when cells were grown in medium with Tween 80.

In the present study, when cells were grown in medium with Tween 80, the two acid-resistant derivatives exhibited higher mean fatty acid chain length (due to more C18:1 and cycC19:0 content) and lower SFA/UFA ratio than their corresponding parental strains (Table 2). The mean fatty acid chain length, cycC19:0, and SFA/UFA ratio are key factors affecting membrane fluidity (Mykytczuk et al. 2007). Higher mean fatty acid chain length could result in lower membrane fluidity since longer chains more easily span the width of the bilayer, contributing to acyl chain packing and making the membrane environment more gel-like (Cao-Hoang et al. 2008; Wu et al. 2012). Moreover, cycC19:0 is a cyclic fatty acid and generally regarded as a tool to reduce membrane fluidity in many bacteria (Grogan and Cronan 1997; To et al. 2011). However, it was reported that lower SFA/UFA ratio was associated with higher membrane fluidity in some bacteria (Álvarez-Ordóñez et al. 2009; Russell 1984). In fact, we found that the two acid-resistant derivatives displayed lower membrane fluidity (higher anisotropy values) than their corresponding parental strains when cells were grown in medium with Tween 80 (Fig. 4). Previous studies on *Lactobacillus casei* (Wu et al. 2012) and *Desulfovibrio piezophilus* (Khelaifia et al. 2011) reported that the higher mean fatty acid chain length was the major factor contributing to the decrease in membrane fluidity, in spite of the lower SFA/UFA ratio. Therefore, it is likely that the mean fatty acid chain length and cyUFpa-

References

- Álvarez-Ordóñez A, Fernández A, López M, Bernardo A (2009) Relationship between membrane fatty acid composition and heat resistance of acid and cold stressed *Salmonella senftenberg* CECT 4384. *Food Microbiol* 26(3):347–353
- Aricha B, Fishov I, Cohen Z, Sikron N, Pesakhov S, Khozin-Goldberg I, Dagan R, Porat N (2004) Differences in membrane fluidity and fatty acid composition between phenotypic variants of *Streptococcus pneumoniae*. *J Bacteriol* 186(14):4638–4644
- Aureli P, Capurso L, Castellazzi AM, Clerici M, Giovannini M, Morelli L, Poli A, Pregliasco F, Salvini F, Zuccotti GV (2011) Probiotics and health: an evidence-based review. *Pharmacol Res* 63(5):366–376
- Ausubel F, Brent R, Kingston R, Moore D, Seidman J, Smith J (2002) Short protocols in molecular biology, 5th edn. Wiley, New York
- Berger B, Moine D, Mansourian R, Arigoni F (2010) HspR mutations are naturally selected in *Bifidobacterium longum* when successive heat shock treatments are applied. *J Bacteriol* 192(1):256–263
- Broadbent JR, Larsen RL, Deibel V, Steele JL (2010) Physiological and transcriptional response of *Lactobacillus casei* ATCC 334 to acid stress. *J Bacteriol* 192(9):2445–2458
- Broadbent JR, Oberg TS, Hughes JE, Ward RE, Brighton C, Welker DL, Steele JL (2014) Influence of polysorbate 80 and cyclopropane fatty acid synthase activity on lactic acid production by *Lactobacillus casei* ATCC 334 at low pH. *J Ind Microbiol Biotechnol* 41(3):545–553
- Brown JL, Ross T, McMeekin TA, Nichols PD (1997) Acid habituation of *Escherichia coli* and the potential role of cyclopropane fatty acids in low pH tolerance. *Int J Food Microbiol* 37(2–3):163–173
- Cao-Hoang L, Marechal P-A, Mai Lê-Thanh M, Gervais P, Waché Y (2008) Fluorescent probes to evaluate the physiological state and activity of microbial biocatalysts: a guide for prokaryotic and eukaryotic investigation. *Biotechnol J* 3(7):890–903
- Chang YY, Cronan JE (1999) Membrane cyclopropane fatty acid content is a major factor in acid resistance of *Escherichia coli*. *Mol Microbiol* 33(2):249–259
- Chu-Ky S, Tourdot-Marechal R, Marechal PA, Guzzo J (2005) Combined cold, acid, ethanol shocks in *Oenococcus oeni*: effects on membrane fluidity and cell viability. *Biochim Biophys Acta* 1717(2):118–124
- Collado MC, Sanz Y (2007) Induction of acid resistance in *Bifidobacterium*: a mechanism for improving desirable traits of potentially probiotic strains. *J Appl Microbiol* 103(4):1147–1157
- Corcoran BM, Stanton C, Fitzgerald G, Ross RP (2008) Life under stress: the probiotic stress response and how it may be manipulated. *Curr Pharm Des* 14(14):1382–1399
- Cotter PD, Hill C (2003) Surviving the acid test: responses of gram-positive bacteria to low pH. *Microbiol Mol Biol Rev* 67(3):429–453
- Da Silveira MG, Golovina EA, Hoekstra FA, Rombouts FM, Abee T (2003) Membrane fluidity adjustments in ethanol-stressed *Oenococcus oeni* cells. *Appl Environ Microbiol* 69(10):5826–5832
- FAO/WHO (2001) Health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. Report of a joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. Córdoba, Argentina. Available at: <ftp://ftp.fao.org/docrep/fao/009/a0512e/a0512e00.pdf>. Accessed 19 Nov 2004
- Fozo EM, Quivey RG Jr (2004) The *fabM* gene product of *Streptococcus mutans* is responsible for the synthesis of monounsaturated fatty acids and is necessary for survival at low pH. *J Bacteriol* 186(13):4152–4158
- Grogan DW, Cronan JE (1997) Cyclopropane ring formation in membrane lipids of bacteria. *Microbiol Mol Biol Rev* 61(4):429–441
- Gueimonde M, Tölkö S, Korpimäki T, Salminen S (2004) New real-time quantitative PCR procedure for quantification of bifidobacteria in human fecal samples. *Appl Environ Microbiol* 70(7):4165–4169
- Guerzoni ME, Lanciotti R, Cocconcelli PS (2001) Alteration in cellular fatty acid composition as a response to salt, acid, oxidative and thermal stresses in *Lactobacillus helveticus*. *Microbiology* 147(8):2255–2264
- Harmsen HJM, Wildeboer-Veloo ACM, Raangs GC, Wagendorp AA, Klijn N, Bindels JG, Welling GW (2000) Analysis of intestinal flora development in breast-fed and formula-fed infants by using molecular identification and detection methods. *J Pediatr Gastroenterol Nutr* 30(1):61–67
- Jin J, Zhang B, Guo H, Cui J, Jiang L, Song S, Sun M, Ren F (2012) Mechanism analysis of acid tolerance response of *Bifidobacterium longum* subsp. *longum* BBMN 68 by gene expression profile using RNA-sequencing. *PLoS One* 7:e50777. doi:10.1371/journal.pone.0050777
- Johnsson T, Nikkila P, Toivonen L, Rosenqvist H, Laakso S (1995) Cellular fatty acid profiles of *Lactobacillus* and *Lactococcus* strains in relation to the oleic acid content of the cultivation medium. *Appl Environ Microbiol* 61(12):4497–4499
- Kanjee U, Houry WA (2013) Mechanisms of acid resistance in *Escherichia coli*. *Annu Rev Microbiol* 67:65–81
- Khelaifia F, Fardeau ML, Pradel N, Aouissargues C, Garel M, Tamburini C, Cayol JL, Gaudron S, Gail F, Ollivier B (2011) *Desulfovibrio piezophilus* sp. nov., a piezophilic, sulfate-reducing bacterium isolated from wood falls in the Mediterranean Sea. *Int J Syst Evol Microbiol* 61(11):2706–2711
- Kim BH, Kim S, Kim HG, Lee J, Lee IS, Park YK (2005) The formation of cyclopropane fatty acids in *Salmonella enterica* serovar *typhimurium*. *Microbiology* 151(1):209–218
- Langendijk PS, Schut F, Jansen GJ, Raangs GC, Kamphuis GR, Wilkinson MH, Welling GW (1995) Quantitative fluorescence in situ hybridization of *Bifidobacterium* spp. with genus-specific 16S rRNA-targeted probes and its application in fecal samples. *Appl Environ Microbiol* 61(8):3069–3075
- Lay C, Rigottier-Gois L, Holmström K, Rajilic M, Vaughan EE, De Vos WM, Collins MD, Thiel R, Namsolleck P, Blaut M, Doré J (2005) Colonic microbiota signatures across five northern European countries. *Appl Environ Microbiol* 71(7):4153–4155
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. *Methods* 25(4):402–408
- Marteau P, Rambaud JC (1993) Potential of using lactic acid bacteria for therapy and immunomodulation in man. *FEMS Microbiol Rev* 12:207–220
- Matsumoto M, Ohishi H, Benno Y (2004) H^+ -ATPase activity in *Bifidobacterium* with special reference to acid tolerance. *Int J Food Microbiol* 93(1):109–113
- Maus JE, Ingham SC (2003) Employment of stressful conditions during culture production to enhance subsequent cold- and acid-tolerance of bifidobacteria. *J Appl Microbiol* 95(1):146–154
- Mykyteczuk NCS, Trevors JT, Leduc LG, Ferroni GD (2007) Fluorescence polarization in studies of bacterial cytoplasmic membrane fluidity under environmental stress. *Prog Biophys Mol Biol* 95(1–3):60–82
- Oberg TS, Ward RE, Steele JL, Broadbent JR (2013) Genetic and physiological responses of *Bifidobacterium animalis* subsp. *lactis* to hydrogen peroxide stress. *J Bacteriol* 195(16):3743–3751
- Parvez S, Malik KA, Ah Kang S, Kim HY (2006) Probiotics and their fermented food products are beneficial for health. *J Appl Microbiol* 100(6):1171–1185
- Revilla-Guarinos A, Alcántara C, Rozès N, Voigt B, Zúñiga M (2014) Characterization of the response to low pH of *Lactobacillus casei* Δ RR12, a mutant strain with low D-alanylation activity and sensitivity to low pH. *J Appl Microbiol* 116(5):1250–1261
- Ruiz L, Ruas-Madiedo P, Gueimonde M, de los Reyes-Gavilán CG, Margolles A, Sánchez B (2011) How do bifidobacteria counteract

- environmental challenges? Mechanisms involved and physiological consequences. *Genes Nutr* 6(3):307–318
- Russell DA, Ross RP, Fitzgerald GF, Stanton C (2011) Metabolic activities and probiotic potential of bifidobacteria. *Int J Food Microbiol* 149(1):88–105
- Russell NJ (1984) Mechanisms of thermal adaptation in bacteria: blueprints for survival. *Trends Biochem Sci* 9(3):108–112
- Sánchez B, Champomier-Vergès MC, Collado MC, Anglade P, Baraige F, Sanz Y, de los Reyes-Gavilán CG, Margolles A, Zagorec M (2007) Low-pH adaptation and the acid tolerance response of *Bifidobacterium longum* biotype longum. *Appl Environ Microbiol* 73(20):6450–6459
- Sánchez B, de los Reyes-Gavilán CG, Margolles A (2006) The F_1F_0 -ATPase of *Bifidobacterium animalis* is involved in bile tolerance. *Environ Microbiol* 8(10):1825–1833
- Saarela M, Rantala M, Hallamaa K, Nohynek L, Virkajärvi I, Mättö J (2004) Stationary-phase acid and heat treatments for improvement of the viability of probiotic lactobacilli and bifidobacteria. *J Appl Microbiol* 96(6):1205–1214
- Shinitzky M (1984) Membrane fluidity and cellular functions. In: Shinitzky M (ed) *Physiology of membrane fluidity*. CRC Press, Boca Raton, Fla, pp 1–52
- To TMH, Grandvalet C, Tourdot-Maréchal R (2011) Cyclopropanation of membrane unsaturated fatty acids is not essential to the acid stress response of *Lactococcus lactis* subsp. *cremoris*. *Appl Environ Microbiol* 77(10):3327–3334
- Vaugien L, Prevots F, Roques C (2002) Bifidobacteria identification based on 16S rRNA and pyruvate kinase partial gene sequence analysis. *Anaerobe* 8(6):341–344
- Ventura M, Canchaya C, van Sinderen D, Fitzgerald GF, Zink R (2004a) *Bifidobacterium lactis* DSM 10140: identification of the *atp* (*atpBEFHAGDC*) operon and analysis of its genetic structure, characteristics, and phylogeny. *Appl Environ Microbiol* 70(5):3110–3121
- Ventura M, van Sinderen D, Fitzgerald GF, Zink R (2004b) Insights into the taxonomy, genetics and physiology of bifidobacteria. *Antonine Van Leeuwenhoek* 86:205–223
- Vitali B, Turrone S, Serina S, Sosio M, Vannini L, Candela M, Guerzoni ME, Brigidi P (2008) Molecular and phenotypic traits of in-vitro-selected mutants of *Bifidobacterium* resistant to rifaximin. *Int J Antimicrob Agents* 31(6):555–560
- Waddington L, Cyr T, Hefford M, Hansen LT, Kalmokoff M (2010) Understanding the acid tolerance response of bifidobacteria. *J Appl Microbiol* 108(4):1408–1420
- Wang Y, Corrieu G, Béal C (2005) Fermentation pH and temperature influence the cryotolerance of *Lactobacillus acidophilus* RD758. *J Dairy Sci* 88(1):21–29
- Wu C, Huang J, Zhou R (2014) Progress in engineering acid stress resistance of lactic acid bacteria. *Appl Microbiol Biotechnol* 98(3):1055–1063
- Wu C, Zhang J, Wang M, Du G, Chen J (2012) *Lactobacillus casei* combats acid stress by maintaining cell membrane functionality. *J Ind Microbiol Biotechnol* 39(7):1031–1039
- Yang Y, Kadim MI, Khoo WJ, Zheng Q, Setyawati MI, Shin Y-J, Lee S-C, Yuk H-G (2014a) Membrane lipid composition and stress/virulence related gene expression of *Salmonella enteritidis* cells adapted to lactic acid and trisodium phosphate and their resistance to lethal heat and acid stress. *Int J Food Microbiol* 191:24–31
- Yang Y, Khoo WJ, Zheng Q, Chung H, Yuk H-G (2014b) Growth temperature alters *Salmonella enteritidis* heat/acid resistance, membrane lipid composition and stress/virulence related gene expression. *Int J Food Microbiol* 172:102–109
- Yuk H-G, Marshall DL (2004) Adaptation of *Escherichia coli* O157: H7 to pH alters membrane lipid composition, verotoxin secretion, and resistance to simulated gastric fluid acid. *Appl Environ Microbiol* 70(6):3500–3505