

RESEARCH PAPER

# Identification of conserved and novel microRNAs that are responsive to heat stress in *Brassica rapa*

Xiang Yu<sup>1</sup>, Han Wang<sup>1</sup>, Yizhen Lu<sup>2</sup>, Marjo de Ruiter<sup>3</sup>, Mike Cariaso<sup>3</sup>, Marcel Prins<sup>3</sup>, Arjen van Tunen<sup>3</sup> and Yuke He<sup>1,\*</sup>

<sup>1</sup> National Key Laboratory of Plant Molecular Genetics, Shanghai Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, 300 Fenglin Road, Shanghai 200032, China

<sup>2</sup> James D. Watson Institute of Genome Sciences, Zhejiang University, 268 Kaixuan Road, Hangzhou 310012, China

<sup>3</sup> Keygene, N.V., Agro Business Park 90, 6708 PW Wageningen, The Netherlands

\* To whom correspondence should be addressed. E-mail: [ykhe@sibs.ac.cn](mailto:ykhe@sibs.ac.cn)

Received 1 August 2011; Revised 22 September 2011; Accepted 27 September 2011

## Abstract

The species *Brassica rapa* includes various vegetable crops. Production of these vegetable crops is usually impaired by heat stress. Some microRNAs (miRNAs) in *Arabidopsis* have been considered to mediate gene silencing in plant response to abiotic stress. However, it remains unknown whether or what miRNAs play a role in heat resistance of *B. rapa*. To identify genomewide conserved and novel miRNAs that are responsive to heat stress in *B. rapa*, we defined temperature thresholds of non-heading Chinese cabbage (*B. rapa* ssp. *chinensis*) and constructed small RNA libraries from the seedlings that had been exposed to high temperature (46 °C) for 1 h. By deep sequencing and data analysis, we selected a series of conserved and novel miRNAs that responded to heat stress. In total, Chinese cabbage shares at least 35 conserved miRNA families with *Arabidopsis thaliana*. Among them, five miRNA families were responsive to heat stress. Northern hybridization and real-time PCR showed that the conserved miRNAs bra-miR398a and bra-miR398b were heat-inhibitive and guided heat response of their target gene, *BracCSD1*; and bra-miR156h and bra-miR156g were heat-induced and its putative target *BracSPL2* was down-regulated. According to the criteria of miRNA and miRNA\* that form a duplex, 21 novel miRNAs belonging to 19 miRNA families were predicted. Of these, four were identified to be heat-responsive by Northern blotting and/or expression analysis of the putative targets. The two novel miRNAs bra-miR1885b.3 and bra-miR5718 negatively regulated their putative target genes. 5'-Rapid amplification of cDNA ends PCR indicated that three novel miRNAs cleaved the transcripts of their target genes where their precursors may have evolved from. These results broaden our perspective on the important role of miRNA in plant responses to heat.

**Key words:** *Brassica rapa*, heat response, miRNA, small RNA.

## Introduction

The species *Brassica rapa* includes various vegetable crops. Comparative genomic study reveals the conserved linkage arrangements and collinear chromosome segments between *B. rapa* and *Arabidopsis thaliana*, which diverged from a common ancestor approximately 13–17 million years ago (Mun *et al.*, 2009). The *B. rapa* genome contains triplicated homologous counterparts of corresponding segments of the *A. thaliana* genome due to triplication of the entire

genome (whole-genome triplication). Production of these vegetable crops is usually impaired by heat stress in many regions. Worldwide, extensive agricultural losses are attributed to heat, often in combination with drought or other stresses (Mittler, 2006). Although heat-resistant molecular breeding has been possible, genetic improvement of crops is hindered by a lack of gene resources relating to heat resistance.

Abbreviations: miRNA, microRNA; RACE, rapid amplification of cDNA ends; siRNA, small interfering RNA; ta-siRNA, *trans*-acting small interfering RNA. TE, transposable element; EST, expressed sequence tag; SSC, 0.15 M NaCl/0.015 M sodium citrate; RT, reverse transcriptase; nt, nucleotide; NT, normal temperature; HT, high temperature.

© 2011 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Plant endogenous small non-coding RNAs are involved in the plant response to abiotic stress, and are divided into four categories: microRNAs (miRNAs), *trans*-acting small interfering RNAs (ta-siRNAs), natural antisense transcripts siRNAs, and repeat associated siRNAs (Jamalkandi and Masoudi-Nejad, 2009). Plant canonical miRNAs are  $\approx$ 21-nucleotide (nt) small RNAs that are processed by the DCL1/HYL1 complex from stem-loop precursors (Kurihara *et al.*, 2006), and are loaded to the RISC complex for regulating plant development and stress response (Shukla *et al.*, 2008; Chen, 2009). Recently, a more complex mechanism of miRNA processing in plants and animals was identified. Most known miRNAs are derived from intergenic regions, while some of them are produced from introns (called mirtrons) (Babiarz *et al.*, 2008; Zhu *et al.*, 2008), exons (Li *et al.*, 2010a), transposable elements (TEs) (Piriyaopongsa *et al.*, 2008; Devor *et al.*, 2009), and even tRNA (Miyoshi *et al.*, 2010). Meanwhile, more evidence for crosstalk between the miRNA pathways and other siRNA pathways has also been reported. For example, the secondary siRNAs are triggered by 22-nt miRNA rather than by canonical 21-nt miRNA (Axtell *et al.*, 2006; Cuperus *et al.*, 2010; Chen *et al.*, 2010). The miRNA hairpins are also processed by the DCL3/RDR2 pathway to produce 24-nt siRNA (Vazquez *et al.*, 2008; Chellappan *et al.*, 2010). miRNAs are known to function in cleaving target genes and inhibiting translation (Chen, 2004; Brodersen *et al.*, 2008; Lanet *et al.*, 2009), whereas 24-nt siRNAs from the miRNA precursors are involved in the methylation of the target genes at the transcriptional level (Chellappan *et al.*, 2010; Wu *et al.*, 2010). On the other hand, miRNA-targeted genes have been identified by high-throughput degradome sequencing in *Arabidopsis* and rice (Addo-Quay *et al.*, 2008; German *et al.*, 2008; Li *et al.*, 2010b). miRNAs and their targets play a very important role in the vegetative phase and floral transition (Wang *et al.*, 2009; Wu *et al.*, 2009), hormone biosynthesis and signaling (Mallory *et al.*, 2005; Reyes and Chua, 2007), and polarity formation and morphogenesis (Sieber *et al.*, 2007; Zhou *et al.*, 2007; Liu *et al.*, 2010; Liu *et al.*, 2011). Others function in stress resistance such as drought (Li *et al.*, 2008), over-oxidation (Sunkar *et al.*, 2006), and phosphate starvation (Fujii *et al.*, 2005). One of the miRNAs associated with stress tolerance is miR398, the expression of which is transcriptionally down-regulated by oxidative stress. In *Arabidopsis*, miR398 was found to target the two closely related Cu/Zn superoxide dismutase-coding genes *CSD1* and *CSD2*, and to regulate plant tolerance to oxidative stress conditions (Sunkar *et al.*, 2006). Recently, the genome-wide analysis of miRNAs under heat stress conditions in wheat has been reported (Xin *et al.*, 2010). In general, identification of conserved and novel heat-responsive miRNAs in *B. rapa* could advance our understanding of their functions in plant heat resistance.

Heat stress disturbs cellular homeostasis and can lead to leaf etiolation, severe retardation in growth and development, and even death. The accumulation of heat-shock proteins under the control of heat stress transcription

factors is assumed to play a central role in the heat stress response and in acquired thermotolerance in plants (Kotak *et al.*, 2007). Several proteins, such as DREB2A (Sakuma *et al.*, 2006), MBF1C (Suzuki *et al.*, 2005), and CTL1 (Kwon *et al.*, 2007), have been shown to improve the heat resistance of plants when over-accumulated. Some signal pathways such as ethylene, abscisic acid (Larkindale *et al.*, 2005), hydrogen peroxide, and inositol trisphosphate (Liu *et al.*, 2006) are involved in crosstalk with pathways of plant thermotolerance. How these proteins protect specific critical targets and whether small RNAs regulate these proteins are major open questions.

With the technological development of small RNA deep sequencing, many miRNAs have been discovered in various crops (Zhu *et al.*, 2008; Lelandais-Briere *et al.*, 2009; Pantaleo *et al.*, 2010). The species *B. rapa* is one of the crops that is mostly closely related to the model plant *Arabidopsis thaliana* (Snowdon, 2007). Vegetable crops belonging to *B. rapa* are very sensitive to heat stress. Losses in the yield and quality of these crops occur especially in summer and in warm regions. Recently, great progress has been made in the sequencing and annotating of *B. rapa* genomes, making it possible to conduct a genome-wide survey of miRNAs in *B. rapa*. Identification of heat-responsive miRNAs offers the opportunity to understand mechanisms of plant response to heat stress.

## Materials and methods

### Small RNA deep sequencing

Wu-11, an inbred line belonging to non-heading Chinese cabbage (*B. rapa* ssp. *chinensis*), was used for constructing small RNA libraries. All plants were grown under a 16/8 h light/dark photoperiod at 22 °C for 3 weeks, and some of the samples were treated with heat shock at 46 °C for 1 h. RNA samples from aboveground parts of the seedlings were prepared using the mirVana miRNA Isolation Kit (Ambion, Austin, TX, USA) and the Alternative v1.5 Protocol (Illumina, San Diego, CA, USA), and small RNA sequencing was performed using Illumina GAII sequencer, according to the manufacturer's protocol.

### Analysis of conserved miRNAs

We aligned our small RNA deep-sequencing data with the mature sequences of *Arabidopsis* miRNA families in miRBase, allowing two mismatches. We then matched all potential miRNAs with *Brassica* genomic Scaffold sequence (<http://brassicadb.org>) or expressed sequence tag (EST) databases from the Brassica Genome Gateway (<http://brassica.bbsrc.ac.uk/>) to export 600-bp flanking sequences from their genomic loci or whole EST sequences. Both of these types of sequence were used to predict secondary structures using RNAfold software. Finally, we identified whether these precursors satisfied the standard of miRNA stem-loop structure (Meyers *et al.*, 2008).

### Identification of novel miRNAs in *B. rapa*


All of the small RNAs we obtained from small RNA deep sequencing were blasted with the *Brassica* genomic Scaffold database and *Brassica* ESTs (see above). Small RNAs that had fewer than 10 hits in the genome were retained. From these databases, 600-bp flanking sequences of small RNAs were used to predict their secondary structures using RNAfold software.

Accordingly, a judgement was made by determining whether the small RNAs and their reverse sequences were on the stem with fewer than four bulges (more than three continuous bulges also not been allowed), the standard miRNA stem-loop described in the literature (Meyers *et al.*, 2008). To investigate the origins of the proper precursors of the stem-loop, the small RNAs were aligned with the *Arabidopsis* TAIR9 cDNA database by local BLASTN.

The NT1 and HT1 treatments generated 14.67 million and 12.77 million small RNA reads in the first experiment; and NT2 and HT2 treatments generated 11.25 million and 14.61 million small RNA reads in the second experiment (Table 1). Because of the difference in abundance of total small RNAs between NT and HT data sets in each experiment, we normalized the abundance of each data set to 10 million [units of transcripts per 10 million (TP10M)].

Some small RNAs were seen in the HT data sets rather than in the NT data sets and hence designated as HT1- or HT2-enriched, while others were seen in the NT data sets rather than in the HT data sets and regarded as NT1- or NT2-enriched. HT-enriched small RNAs (28 and 29% for HT1 and HT2) were more abundant than NT-enriched ones (22 and 24% for NT1 and NT2), meaning that heat stress induces some small RNAs and represses others, and that more unique small RNAs are induced than repressed. Nevertheless, the majority of small RNAs were shared by the HT and NT data sets as they were seen in common for the two populations.

Small RNA sequences of Chinese



**Fig. 1.** Length distribution of small RNAs in two databases of non-heading Chinese cabbage after HT and NT treatments (NT1 and HT1, and NT2 and HT2, from experiments 1 and 2, respectively). (A) Length distribution of unique sequences in NT and HT small-RNA data sets. (B) Length distribution of total reads in NT and HT small-RNA data sets.

we did not find miR173 and miR173-derived ta-siRNA sequences or their homologues in all of the four small-RNA data sets. To examine whether there are any homologous genes of *MIR173* in Chinese cabbage, we searched the *Brassica* genomic Scaffold database. No genomic sequences of Chinese cabbage were homologous to the *MIR173*, *TAS1*, or *TAS2* genes in *Arabidopsis*. Probably, the genome of Chinese cabbage is lacking the *MIR173*, *TAS1*, and *TAS2* genes. In contrast, miR390 and the corresponding ta-siRNA sequences were present in our small-RNA data set and were matched to the *Brassica* scaffold database, indicating that the *MIR390* and *TAS3* genes in Chinese cabbage are functional. In addition, the bra-miR828 that targets at *TAS4* was detected in Chinese cabbage but an siRNA generated from *TAS4* was not found. We deduce

that the ta-siRNAs from the *TAS1*, *TAS2*, and *TAS4* genes may not exist in Chinese cabbage. Nevertheless, neither miR390 nor ta-siRNAs from *TAS3* were affected by high temperature.

After annotating the conserved miRNA, we were able to examine which miRNAs are responsive to heat stress according to the differential expression analysis of miRNAs. The 35 conserved miRNA families in Chinese cabbage include 62 sequence-specific miRNAs. Among heat-responsive miRNAs, four were up-regulated and three were down-regulated with a more than 1.5-fold change (Table 3). Remarkably, bra-miR398a and bra-miR398b were down-regulated with more than 10- and 3-fold changes, respectively. bra-miR398a and bra-miR398b are derived from two homologous precursors of miR398, and have one mismatch

**Table 2.** Total members (IM) represent the number of Chinese cabbage bra-miRNA identical to Arabidopsis miRNA families that are highly conserved between the Brassicaceae family and other species.

bra-miRNA similar to ath-miRNA with one or two

represents the miRNA families that are highly conserved


between the Brassicaceae family and other species

Brassicaceae family. L represents the

specific miRNA families.

**Table 3.**

A 5' TGTGTTCTCAGGTCACCCCTG 3' bra-miR398b  
 5' TGTGTTCTCAGGTCACCCCTT 3' bra-miR398a  
 |||| | ||||| | |||||  
 3' A C A C T A G A G T C C C A T G G G G A A 5' *BracCSD1*  
 5' T G A C A G A A G A A G A G A G C A C 3' bra-miR156h  
 5' C G A C A G A A G A G A G T G A G C A C 3' bra-miR156g  
 ||||| ||||| ||||| ||||| |||||  
 3' A C T G T C T T C T C T C T C T G T G 5' *BracSPL2*



**Fig. 2.** Accumulation of bra-miR398a and expression of miR398 and miR156 targets under heat stress. (A) Alignment of the miRNAs with the targets: bra-miR398 with *BracCSD1*

in sequence. The homologue of bra-miR398 in *Arabidopsis* responds to oxidative stress by regulating its target *CSD1*, a Cu/Zn superoxide dismutase (Sunkar *et al.*, 2006). To confirm the heat response of bra-miR398, we performed Northern blotting of miR398 and real-time PCR of the target gene. The accumulation of both bra-miR398a and bra-miR398b in the HT seedlings was much lower than in the NT seedlings (Fig. 2A and B). On the contrary, the expression of *BracCSD1*, as a target gene of bra-miR398 (Supplementary Table S4), was much higher in the HT seedlings than in the NT seedlings (Fig. 2C). This result

indicates that *BracCSD1* regulates heat response of *B. rapa* and that bra-miR398 guides the *BracCSD1* gene by silencing it. miR156h and miR156g were especially up-regulated in contrast to miR156a-f. We validated that

*BracSPL2* was sharply down-regulated under heat stress (Fig. 2D), and therefore designated it as the proper target of miR156h and miR156g in heat response. miR399 is repressed and miR167 is induced by heat stress, consistent with a report in wheat (Xin *et al.*, 2010).

#### *Analysis of heat-responsive small RNAs originating from miRNA precursors*

Many small RNAs are generated from the miRNA precursors in *Arabidopsis*. Some of them, called miRNA-sibling RNAs, also play a post-transcriptional role by cleaving target genes (Zhang *et al.*, 2010). According to the distribution of small RNAs in miRNA precursors, small RNAs are classified into three types: miRNA variants overlapped with miRNA, miRNA\* variants overlapped with miRNA\*, and miRNA-sibling RNAs in the flanking sequences of miRNAs or miRNA\*. Consistent with the heat-responsive miRNAs, some small siRNAs from the same precursor display the same pattern of heat response. For example, both the miRNA variants and miRNA\* variants of bra-miR156h-2 were up-regulated under heat stress (Fig. 3A). However, the heat-responses of a few miRNA\* variants is different from that of their own miRNA variants.

Accumulation of the miRNA\* variants of bra-miR167a and bra-miR400 was repressed by heat stress whereas their miRNA variants were slightly up-regulated or unchanged (Fig. 3B and C). These results suggest that high temperature affects the processing or stability of miRNAs and/or their miRNA\* variants in different ways.

#### *Identification of novel miRNAs responsive to heat stress*

To predict novel miRNAs in *Brassica*, we matched all of the small RNAs (20–22 bp) of Chinese cabbage with the *Brassica* scaffold database and EST. The 600-bp flanking fragments of small RNAs were used to draw their secondary structure using RNAfold. Candidate precursors were selected according to whether the small RNAs and their reverse sequences were on the stem, with fewer than four bulges, according to one of the miRNA standards (Meyers *et al.*, 2008). Then the candidate precursors were aligned with the *Arabidopsis* cDNA database using local BLASTN. Lastly, all small RNAs in our database were matched to the candidate precursors and the distribution of the small RNAs was mapped along the candidate precursors. We excluded the precursors homologous to rRNA, most TE genes, and those with many smear sequences. According to the miRNA/miRNA\* criteria, 21 novel miRNAs belonging to 19 miRNA families were selected (Table 4, Supplementary Table S5). Potential targets of all of the novel miRNA were predicted according to the complementarity between miRNAs and EST sequences (Supplementary Table S6).

Among the novel miRNAs, bra-miR5714 and bra-miR5726 were up-regulated and bra-miR5716 and bra-miR1885b.3 were down-regulated by more than 2-fold under heat stress (Table 3). bra-miR1885b.3 precursors were able to produce

three pairs of miRNA/miRNA\* (Fig. 4A and C). Under heat stress, the accumulations of bra-miR1885b.3, bra-miR1885b.3\*, and bra-miR1885b.2\* were repressed severely (Fig. 4B). To confirm the heat response of these novel miRNAs, we performed Northern blotting of small RNA. The accumulation of bra-miR1885b.3 in HT seedlings was reduced sharply by heat stress (Fig. 4D). In contrast, its putative target gene was up-regulated under the same condition (Fig. 4E). We also tested the expression of the putative targets of bra-miR5714 and bra-miR5726, and found that they were repressed under heat-stress, corresponding to the two miRNAs that were induced by heat stress (Fig. 4F and G).

Another potential heat-responsive novel miRNA is bra-miR5718 (Fig. 5A and B). Under heat stress, the accumulation of bra-miR5718 was increased by more than 1.5-fold. In contrast, *BracPAPI0*, the putative target gene of bra-miR5718, was remarkably down-regulated (Fig. 5C). We noticed that most of the small RNAs originated from the bra-miR5718 precursor were also heat-induced. These results reveal that bra-miR1885b.3 and bra-miR5718 are two novel miRNAs that function in the *B. rapa* heat response.

#### *Evolutionary relationship between novel miRNAs and their targets*

By comparing the sequence similarity of the putative miRNA precursors against the *Arabidopsis* cDNA database by BLASTN, we found that the putative precursors of 10 novel miRNAs were homologous to certain fragments of the protein-coding genes. Among these novel miRNAs, three cleaved their targets, from which they may have evolved. The precursor of bra-miR5713 may evolve from the gene *BracDRL1* (*DEFORMED ROOTS AND LEAVES 1*) (Fig. 6A). As indicated in Fig. 6B, bra-miR5713-3p and bra-miR5713-5p are two small RNAs that are derived from the same precursor. The low-abundance bra-miR5713-3p was predicted to target *BracDRL1* while the high-abundance bra-miR5713-5p was predicted to target *BracVEL1*. To define the cleavage sites of these miRNA-targeted genes, we performed 5'-RACE experiments. As expected, the transcripts of *BracDRL1* and *BracVEL1* were cleaved in the regions complimentary with bra-miR5713-3p and bra-miR5713-5p, respectively. Similarly, bra-miR1885b.1 and bra-miR1885b.3 are two small RNAs from the same precursor. They may have evolved from the gene *BracTAO1* (Fig. 6C). 5'-RACE experiments showed that bra-miR1885b.1 rather than bra-miR1885b.3 cleaves *BracTAO1* (Fig. 6D). We suggest that miRNAs derived from the same miRNA precursors regulate different targets at the post-transcriptional level. The third miRNA that may have evolved from its target was bra-miR5718 (Fig. 5A). Our 5'-RACE PCR confirmed that the transcripts of *BracPAPI0* were cleaved by bra-miR5718 (Fig. 5B).

We noticed that the major cleavage sites in the transcripts of several miRNA targets were not in the center of the binding sites. This indicated the broad variance of cleavage





**Table 4.** The sequences of novel miRNAs in *B. rapa*.

Novel miRNA	Sequence	Length (nt)	Locus number
bra-miR1140	ACAGCCTAAACCAATCGGAGC	21	1
bra-miR1885.2	TACATCTTCTCCGCGGAAGCTC	22	2
bra-miR1885b.3	ATTGTGGACAAAGAAGGAAG	20	1
bra-miR5654	ATAAATCCCAAGCATCATCCA	21	2
bra-miR5711	TGTTTTGTGGGTTTCTACCGA	22	1
bra-miR5712	AATATTAATAATTGGTGAG	21	1
bra-miR5713-3p	TTCTAAGCCTACCTAGCTCCGG	22	1
bra-miR5713-5p	AGGCTTAGAAGAACGTTTGTT	21	1
bra-miR5714	AGACTCTACGACATCAAGAAAC	22	1
bra-miR5715	ACGTGATAAGCCTCTGAAGAA	21	1
bra-miR5716	TTGGATAATTGAAGATATAAA	21	1
bra-miR5717	GTTTGGATTGTTGCCTTGGC	21	1
bra-miR5718	TCAGAACCAAACACAGAACAAG	22	1
bra-miR5719	TTGTGATGATAATACGACTTC	21	1
bra-miR5720	TTGTGATTTGGTTGGAATATC	21	1
bra-miR5721	AAAAATGGAGTGAGAAATGGA	21	1
bra-miR5722	TGAAATAGAGTCATGTGGAACG	22	1
bra-miR5723	AATGTGCTGCAATATCTCTGC	21	1
bra-miR5724	AACCGCCGGTTTGATAATAGC	21	1
bra-miR5725	ATTTGGCACAATCTGATCTGC	21	1
bra-miR5726	CAAAGGTTGCTTGAATAAGGT	21	1

genes. By comparing the known miRNAs and the miRNA precursors, we identified 35 miRNA families conserved between Chinese cabbage and *Arabidopsis*. The conserved miRNAs in Chinese cabbage are identical or similar (with one or two mismatches) to their counterparts in *Arabidopsis*. *Brassica* species are closely related to the model plant *A. thaliana*. The gene order within their genomes has remained extensively collinear and remarkably conserved over a long evolutionary period. Although Chinese cabbage is diploid, its genome shows characteristics of triplication (Town *et al.*, 2006; Mun *et al.*, 2009). Many *MIRNA* genes in *B. rapa* show far more copies than in *Arabidopsis*. In *B. rapa*, mature miRNA sequences are more diverse than those of *Arabidopsis*. For example, the precursor of bra-miR156d-2 shares the most homology with ath-miR156d among all members of the miR156 family, but the mature miRNA contains one nucleotide substitution compared with ath-miR156d (Supplementary Table S2). In total, 22 miRNAs from the conserved miRNA families in *Brassica* show one or two nucleotide substitutions compared with those of *Arabidopsis* (Table 2).


Recently, many types of miRNAs have been identified in the genome of *A. lyrata*, in which 13% are species-specific, because they are not found in its close relative *A. thaliana* (Zhang *et al.*, 2010). Only one *Brassica*-specific miRNA has been reported (He *et al.*, 2008). Using the almost complete genomic sequences of *B. rapa*, we selected 19 novel miRNA families according to the strict miRNA/miRNA\* criterion. Interestingly, some novel miRNA precursors have evolved from their target genes. For example, bra-miR5713, bra-miR1885, and bra-miR5718 are evolved from their targets *BracDRL1*, *BracTAO1*, and *BracPAP10*, respectively. Kutter *et al.* (2007) reported that miR824 was duplicated from its target *AGL16*. We suggest that co-evolution

between miRNA genes and their target genes is partly responsible for the diversity of gene silencing.

#### *miRNA is involved in gene-regulation pathways of the plant heat response*

A miRNA directs cleavage of a highly complementary target mRNA. Upon heat stress, any change in the type or amount of miRNAs in plant tissues may alter the expression levels of target genes. In this study, the conserved miRNA miR398a is heat-inhibitive while *BracCSD1* gene is heat-sensitive, showing an inverse relationship between miRNA and target. Recently, some components in thermotolerance are found to have the functions in protection from oxidative damage. For example, genetic engineering of the biosynthesis of glycinebetaine enhances thermotolerance of photosystem II in tobacco plants (Yang *et al.*, 2007). Chloroplast NAD(P)H dehydrogenase in tobacco leaves functions in alleviation of oxidative damage caused by temperature stress (Wang *et al.*, 2006). In this study, we found that bra-miR398 and its target genes *BracCSD1* and *BracCSD2* showed an inverse response to heat stress in *B. rapa*. *CSD1* and *CSD2* genes that are the targets of miR398 regulate the plant tolerance to oxidative stress conditions (Sunkar *et al.*, 2006). We wonder whether bra-miR398-mediated *BracCSD1* and *BracCSD2* genes play roles in thermotolerance through protection from oxidative damage.


The miR399 is responsive to phosphate starvation (Fuji *et al.*, 2005). Both of their targets encode putative ubiquitin-conjugating enzyme (UBCs). Under heat stress plant would consume more nitrogen and phosphate; repressing these two miRNAs would accumulate their target genes to adapt the nutrient stress. The miR156s were found to be up-regulated under heat stress in wheat recently (Xin



**Fig. 4.** The inverse expression pattern between bra-miR1885b and the putative target genes under heat stress. (A) Three pairs of miRNA/miRNA\* in the loop hairpin of the bra-miR1885b precursor. (B) Abundance of the miRNAs and miRNA\* derived from the bra-miR1885b precursor under heat stress in the four small-RNA data sets. (C) Alignment of three novel miRNAs with the binding sites of their predicted targets. (D) Northern blotting of miR1885b.3 under heat stress. (E) RT-PCR showing relative expression of the putative target genes under heat stress. (F) Real-time PCR showing relative expression of miR5714-targeted AM38896 (AM38896). (G) Real-time PCR showing relative expression of miR5726-targeted EE405869.

*et al.*, 2010), and we found that miR156h and miR156g were specifically induced among the miR156 families. The targets of miR156 are the SPL transcription factor families, which take part in the vegetative phase and floral transition

(Wang *et al.*, 2009; Wu *et al.*, 2009). Among the miR156 targets, *SPL2* were specifically down-regulated under heat stress. The negative correlation between miR156h/miR156g and *SPL2* may suggest that miR156h and miR156g may



**Fig. 5.** Evolutionary relationship and heat-response of bra-MIR5718 and its target *BracPAP10*. (A) Heat-responsive small RNAs originated from bra-miR5718 precursors. Arabic numbers after the sequences are the normalized reads of small RNAs in the NT1, HT1, NT2, and HT2 data sets, respectively. (B) Alignment of bra-miR5718 precursors with *BracPAP10*. The reverse sequence and direction of bra-miR5718 are indicated with a reverse arrow, while the DNA sequence of *BracPAP10* is underlined with a forward arrow. (C) Northern blotting bra-miR5718 under heat stress. (D) RT-PCR showing the expression of the target *BracPAP10* under heat stress. (E) The position of dominant 5'-RACE products of *BracPAP10* mRNA is indicated by a vertical arrow in the expanded region.

target *SPL2* effectively rather than *SPL3*, *SPL9*, and *SPL10*, which are regulated by miR156a. However, the downstream targets of *SPL2* are still unknown.

The novel miRNAs bra-miR1885b.3 and bra-miR5718 are responsive to heat stress, in contrast to their targets. To identify function of bra-miR398, bra-miR1885b.3, and bra-miR5718 in terms of thermotolerance, an attempt has been made in our laboratory to construct transgenic plants overexpressing these miRNAs and their target genes. Identification of the conserved and novel miRNAs involved in the heat response of *B. rapa* will facilitate our un-

derstanding of the molecular mechanisms governing the plant heat response and thermotolerance.

## Supplementary material

Supplementary material is available at *JXB* online.

Supplementary Fig. S1. Number of miRNAs belonging to 33 families conserved among 14 plant species the whole-genome sequences of which are available in miRBase.

Supplementary Table S1. Primers used in 5'-RACE PCR and real-time PCR.



## Acknowledgements

This work was supported by grants from the Natural Science Foundation of China (30730053) and 973 Program of China (2012CB113903).

## References

- Addo-Quaye C, Eshoo TW, Bartel DP, Axtell MJ.** 2008. Endogenous siRNA and miRNA targets identified by sequencing of the *Arabidopsis* degradome. *Current Biology* **18**, 758–762.
- Axtell MJ, Jan C, Rajagopalan R, Bartel DP.** 2006. A two-hit trigger for siRNA biogenesis in plants. *Cell* **127**, 565–577.
- Babiarz JE, Ruby JG, Wang Y, Bartel DP, Blelloch R.** 2008. Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes and Development* **22**, 2773–2785.
- Brodersen P, Sakvarelidze-Achard L, Bruun-Rasmussen M, Dunoyer P, Yamamoto YY, Sieburth L, Voinnet O.** 2008. Widespread translational inhibition by plant miRNAs and siRNAs. *Science* **320**, 1185–1190.
- Chellappan P, Xia J, Zhou X, Gao S, Zhang X, Coutino G, Vazquez F, Zhang W, Jin H.** 2010. siRNAs from miRNA sites mediate DNA methylation of target genes. *Nucleic Acids Research* **38**, 6883–6894.
- Chen HM, Chen LT, Patel K, Li YH, Baulcombe DC, Wu SH.** 2010. 22-Nucleotide RNAs trigger secondary siRNA biogenesis in plants. *Proceedings of the National Academy of Sciences, USA* **107**, 15269–15274.
- Chen K, Rajewsky N.** 2007. The evolution of gene regulation by transcription factors and microRNAs. *Nature Reviews Genetics* **8**, 93–103.
- Chen X.** 2004. A MicroRNA as a translational repressor of *APETALA2* in *Arabidopsis* flower development. *Science* **303**, 2022–2024.
- Chen X.** 2009. Small RNAs and their roles in plant development. *Annual Review of Cell and Developmental Biology* **25**, 21–44.
- Cuperus JT, Carbonell A, Fahlgren N, Garcia-Ruiz H, Burke RT, Takeda A, Sullivan CM, Gilbert SD, Montgomery TA, Carrington JC.** 2010. Unique functionality of 22-nt miRNAs in triggering RDR6-dependent siRNA biogenesis from target transcripts in *Arabidopsis*. *Nature Structural and Molecular Biology* **17**, 997–1003.
- de Felippes FF, Schneeberger K, Dezulian T, Huson DH, Weigel D.** 2008. Evolution of *Arabidopsis thaliana* microRNAs from random sequences. *RNA* **14**, 2455–2459.
- Devor EJ, Peek AS, Lanier W, Samollow PB.** 2009. Marsupial-specific microRNAs evolved from marsupial-specific transposable elements. *Gene* **448**, 187–191.
- Fahlgren N, Jogdeo S, Kasschau KD, Sullivan CM, Chapman EJ, Laubinger S, Smith LM, Dasenko M, Givan SA, Weigel D, Carrington JC.** 2010. MicroRNA gene evolution in *Arabidopsis lyrata* and *Arabidopsis thaliana*. *The Plant Cell* **22**, 1074–1089.
- Fujii H, Chiou TJ, Lin SI, Aung K, Zhu JK.** 2005. A miRNA involved in phosphate-starvation response in *Arabidopsis*. *Current Biology* **15**, 2038–2043.
- German MA, Pillay M, Jeong DH, et al.** 2008. Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. *Nat Biotechnol* **26**, 941–946.
- He XF, Fang YY, Feng L, Guo HS.** 2008. Characterization of conserved and novel microRNAs and their targets, including a TuMV-induced TIR-NBS-LRR class R gene-derived novel miRNA in *Brassica*. *FEBS Letters* **582**, 2445–2452.
- Hsieh LC, Lin SI, Shih AC, Chen JW, Lin WY, Tseng CY, Li WH, Chiou TJ.** 2009. Uncovering small RNA-mediated responses to phosphate deficiency in *Arabidopsis* by deep sequencing. *Plant Physiology* **151**, 2120–2132.
- Jamalkandi SA, Masoudi-Nejad A.** 2009. Reconstruction of *Arabidopsis thaliana* fully integrated small RNA pathway. *Functional and Integrated Genomics* **9**, 419–432.
- Jones-Rhoades MW, Bartel DP, Bartel B.** 2006. MicroRNAs and their regulatory roles in plants. *Annual Review of Plant Biology* **57**, 19–53.
- Kotak S, Larkindale J, Lee U, von Koskull-Doring P, Vierling E, Scharf KD.** 2007. Complexity of the heat stress response in plants. *Current Opinion in Plant Biology* **10**, 310–316.
- Kurihara Y, Takashi Y, Watanabe Y.** 2006. The interaction between DCL1 and HYL1 is important for efficient and precise processing of pri-miRNA in plant microRNA biogenesis. *RNA* **12**, 206–212.
- Kutter C, Schob H, Stadler M, Meins F, Jr., Si-Ammour A.** 2007. MicroRNA-mediated regulation of stomatal development in *Arabidopsis*. *The Plant Cell* **19**, 2417–2429.
- Kwon Y, Kim SH, Jung MS, Kim MS, Oh JE, Ju HW, Kim KI, Vierling E, Lee H, Hong SW.** 2007. *Arabidopsis* hot2 encodes an endochitinase-like protein that is essential for tolerance to heat, salt and drought stresses. *The Plant Journal* **49**, 184–193.
- Lanet E, Delannoy E, Sormani R, Floris M, Brodersen P, Crete P, Voinnet O, Robaglia C.** 2009. Biochemical evidence for translational repression by *Arabidopsis* microRNAs. *The Plant Cell* **21**, 1762–1768.
- Larkindale J, Hall JD, Knight MR, Vierling E.** 2005. Heat stress phenotypes of *Arabidopsis* mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiology* **138**, 882–897.
- Lelandais-Briere C, Naya L, Sallet E, Calenge F, Frugier F, Hartmann C, Gouzy J, Crespi M.** 2009. Genome-wide *Medicago truncatula* small RNA analysis revealed novel microRNAs and isoforms differentially regulated in roots and nodules. *The Plant Cell* **21**, 2780–2796.
- Li T, Li H, Zhang YX, Liu JY.** 2010a. Identification and analysis of seven H<sub>2</sub>O<sub>2</sub>-responsive miRNAs and 32 new miRNAs in the seedlings of rice (*Oryza sativa* L. ssp. *indica*). *Nucleic Acids Research* **39**, 2821–2833.
- Li WX, Oono Y, Zhu J, He XJ, Wu JM, Iida K, Lu XY, Cui X, Jin H, Zhu JK.** 2008. The *Arabidopsis* NFYA5 transcription factor is regulated transcriptionally and posttranscriptionally to promote drought resistance. *The Plant Cell* **20**, 2238–2251.
- Li YF, Zheng Y, Addo-Quaye C, Zhang L, Saini A, Jagadeeswaran G, Axtell MJ, Zhang W, Sunkar R.** 2010b. Transcriptome-wide identification of microRNA targets in rice. *The Plant Journal* **62**, 742–759.

- Liu HT, Gao F, Cui SJ, Han JL, Sun DY, Zhou RG. 2006. Primary evidence for involvement of IP3 in heat-shock signal transduction in *Arabidopsis*. *Cell Res* **16**, 394–400.
- Liu Z, Jia L, Mao Y, He Y. 2010. Classification and quantification of leaf curvature. *Journal of Experimental Botany* **61**, 2757–2767.
- Liu Z, Jia L, Wang H, He Y. 2011. HYL1 regulates the balance between adaxial and abaxial identity for leaf flattening via miRNA-mediated pathways. *Journal of Experimental Botany* **62**, 4367–4381.
- Mallory AC, Bartel DP, Bartel B. 2005. MicroRNA-directed regulation of *Arabidopsis* AUXIN RESPONSE FACTOR17 is essential for proper development and modulates expression of early auxin response genes. *The Plant Cell* **17**, 1360–1375.
- Meyers BC, Axtell MJ, Bartel B, *et al.* 2008. Criteria for annotation of plant MicroRNAs. *The Plant Cell* **20**, 3186–3190.
- Mittler R. 2006. Abiotic stress, the field environment and stress combination. *Trends Plant Sci* **11**, 15–19.
- Miyoshi K, Miyoshi T, Siomi H. 2010. Many ways to generate microRNA-like small RNAs: non-canonical pathways for microRNA production. *Molecular Genetics and Genomics* **284**, 95–103.
- Mun JH, Kwon SJ, Yang TJ, *et al.* 2009. Genome-wide comparative analysis of the *Brassica rapa* gene space reveals genome shrinkage and differential loss of duplicated genes after whole genome triplication. *Genome Biology* **10**, R111.
- Pantaleo V, Szittyá G, Moxon S, Miozzi L, Moulton V, Dalmay T, Burgyan J. 2010. Identification of grapevine microRNAs and their targets using high-throughput sequencing and degradome analysis. *The Plant Journal* **62**, 960–976.
- Piriyapongsa J, Jordan IK. 2008. Dual coding of siRNAs and miRNAs by plant transposable elements. *RNA* **14**, 814–821.
- Reyes JL, Chua NH. 2007. ABA induction of miR159 controls transcript levels of two MYB factors during *Arabidopsis* seed germination. *The Plant Journal* **49**, 592–606.
- Sakuma Y, Maruyama K, Qin F, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K. 2006. Dual function of an *Arabidopsis* transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression. *Proceedings of the National Academy of Sciences, USA* **103**, 18822–18827.
- Shukla LI, Chinnusamy V, Sunkar R. 2008. The role of microRNAs and other endogenous small RNAs in plant stress responses. *Biochimica Biophysica Acta* **1779**, 743–748.
- Sieber P, Wellmer F, Gheyselinck J, Riechmann JL, Meyerowitz EM. 2007. Redundancy and specialization among plant microRNAs: role of the MIR164 family in developmental robustness. *Development* **134**, 1051–1060.
- Snowdon RJ. 2007. Cytogenetics and genome analysis in *Brassica* crops. *Chromosome Research* **15**, 85–95.
- Sunkar R, Kapoor A, Zhu JK. 2006. Posttranscriptional induction of two Cu/Zn superoxide dismutase genes in *Arabidopsis* is mediated by downregulation of miR398 and important for oxidative stress tolerance. *The Plant Cell* **18**, 2051–2065.
- Suzuki N, Rizhsky L, Liang H, Shuman J, Shulaev V, Mittler R. 2005. Enhanced tolerance to environmental stress in transgenic plants expressing the transcriptional coactivator multiprotein bridging factor 1c. *Plant Physiology* **139**, 1313–1322.
- Town CD, Cheung F, Maiti R, *et al.* 2006. Comparative genomics of *Brassica oleracea* and *Arabidopsis thaliana* reveal gene loss, fragmentation, and dispersal after polyploidy. *The Plant Cell* **18**, 1348–1359.
- Vazquez F, Blevins T, Ailhas J, Boller T, Meins F. Jr 2008. Evolution of *Arabidopsis* MIR genes generates novel microRNA classes. *Nucleic Acids Research* **36**, 6429–6438.
- Wang JW, Czech B, Weigel D. 2009. MiR156-regulated SPL transcription factors define an endogenous flowering pathway in *Arabidopsis thaliana*. *Cell* **138**, 738–749.
- Wang L, Yu X, Wang H, Lu YZ, de Ruiter M, Prins M, He YK. 2011. A novel class of heat-responsive small RNAs derived from the chloroplast genome of Chinese cabbage (*Brassica rapa*). *BMC Genomics* **12**, 289.
- Wang P, Duan W, Takabayashi A, Endo T, Shikanai T, Ye JY, Mi H. 2006. Chloroplastic NAD(P)H dehydrogenase in tobacco leaves functions in alleviation of oxidative damage caused by temperature stress. *Plant Physiology* **141**, 465–474.
- Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS. 2009. The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* **138**, 750–759.
- Wu L, Zhou H, Zhang Q, Zhang J, Ni F, Liu C, Qi Y. 2010. DNA methylation mediated by a microRNA pathway. *Molecular Cell* **38**, 465–475.
- Xin M, Wang Y, Yao Y, Xie C, Peng H, Ni Z, Sun Q. 2010. Diverse set of microRNAs are responsive to powdery mildew infection and heat stress in wheat (*Triticum aestivum* L.). *BMC Plant Biology* **10**, 123.
- Yang X, Wen X, Gong H, Lu Q, Yang Z, Tang Y, Liang Z, Lu C. 2007. Genetic engineering of the biosynthesis of glycinebetaine enhances thermotolerance of photosystem II in tobacco plants. *Planta* **225**, 719–733.
- Zhang W, Gao S, Zhou X, Xia J, Chellappan P, Zhang X, Jin H. 2010. Multiple distinct small RNAs originate from the same microRNA precursors. *Genome Biology* **11**, R81.
- Zhou GK, Kubo M, Zhong R, Demura T, Ye ZH. 2007. Overexpression of miR165 affects apical meristem formation, organ polarity establishment and vascular development in *Arabidopsis*. *The Plant Cell Physiol* **48**, 391–404.
- Zhu QH, Spriggs A, Matthew L, Fan L, Kennedy G, Gubler F, Helliwell C. 2008. A diverse set of microRNAs and microRNA-like small RNAs in developing rice grains. *Genome Research* **18**, 1456–1465.