catalyse initial aromatic ring oxygenation and the elimination of nitrite in the catabolism of nitroaromatic compounds. These enzymes are classified in group III of Rieske non-heme iron dioxygenases (Parales and Resnick, 2006) and are collectively called Nag-like nitroarene dioxygenases. A recently described exception, 3-nitrotoluene dioxygenase (BndA1A2A3A4) is more closely related to biphenyl dioxygenase (Gao et al., 2020).

RW71, which has a broad substrate range including 3,4- and 4,5-dichlorocatechols (Potrawfke et al., 2001). The genes dcnDEF, predicted to encode enzymes that catalyse metabolism of the dichlorocatechol ring-cleavage product, are located at chromosome positions 1 862 862 to 1 865 615. ORFs 10 and 12 are predicted to encode chloromaleylacetate reductase and chlorodienelactone hydrolase. ORF 11 is most closely related to genes encoding chloromuconate cycloisomerase. To further investigate functions of the key genes mentioned above, biochemical characterizations were done to establish the roles of the putative genes in 3,4-DCNB degradation.

DcnAaAbAcAd encodes a dioxygenase converting 3,4-DCNB to 4,5-dichlorocatechol

The position of initial attack on the 3,4-DCNB molecule determines the subsequent downstream catabolic pathway. Attack at the 1,2 position would yield 3,4-dichlorocatechol, whereas attack at the 1,6 position would yield 4,5-dichlorocatechol. In order to establish the catalytic function and the regiospecificity of 3,4-DCNB dioxygenase, we constructed plasmid [pUC18-dcnAaAbAcAd] and transformed it into E. coli strain DH5α to construct E. coli [pUC18-dcnAaAbAcAd] for evaluation of 3,4-DCNB dioxygenase activity via heterologous expression and whole cell biotransformations.

The heterologously expressed nitroarene dioxygenase transformed 3,4-DCNB stoichiometrically to 4,5-dichlorocatechol and nitrite (Fig. 2, Figs S1, S2). The identity of the product was confirmed by HPLC and GC–MS, using commercial 4,5-dichlorocatechol as a standard, and commercial 3,4-dichlorocatechol as a control. The results established that dcnAaAbAcAd encodes a 3,4-DCNB dioxygenase catalysing the conversion of 3,4-DCNB into 4,5-dichlorocatechol with a high degree of



Fig. 2. 3,4-DCNB biotransformation by E.coli DH5α [pUC18-dcnAaAbAcAd]. The disappearance of 3,4-DCNB and the appearance of the products 4,5-dichlorocatechol were quantified by HPLC. The experiments were performed in triplicate, the results shown are the average values from three independent assays, and error bars indicate standard deviations. [Color figure can be viewed at wileyonlinelibrary.com]

Structure modelling of 3,4-DCNB dioxygenase

It is known that the substrate specificity of nitroarene dioxygenase is determined by the large subunit of the terminal oxygenase (Ac subunit) (Parales et al., 1998). The fact that the product of 3,4-DCNB dioxygenation by DcnAcAd is 4,5-dichlorocatechol instead of 3,4-dichlorocatechol indicates that there must be strict control of the orientation of the substrate in the active site. In order to identify the residues that determine the specificity of the oxidation attack, the overall protein structure model of the DcnAc subunit of 3,4-DCNB dioxygenase was built using the 3NT dioxygenase α subunit as template (Fig. 3A). 3,4-DCNB was docked into the substrate pocket of the DcnAc model by Autodock Vina. The mononuclear iron is coordinated by a non-heme iron-binding triad consisting of His206, His211 and Asp360. The substrate 3,4-DCNB binds close to the mononuclear iron, and its aromatic plane faces the ferrous ion with a distance of about 3.9 Å. For substrate binding, some residues, such as Phe200, Val305, Ile207 and Val258, interact with the aromatic ring of 3,4-DCNB through hydrophobic interactions. Tyr222, Gln293 and Asn295 form hydrogen bonds to the nitro group of 3,4-DCNB. A positively charged residue, Lys312, attracts a chlorine atom (4-positon) of 3,4-DCNB with a distance of 4.5 Å (Fig. 3B and C), while the other chlorine atom (3-position) is engaged in halogen–π interaction with the side chain of Trp356 with a distance of 3.1 Å. The key residues mentioned above, together with mononuclear iron atom and water molecules in the active center, place the substrate at a suitable location and orientation. Through analysing this model it can be deduced that C2, remote from catalytic residue Asn295 and close to Val258, cannot be hydroxylated because of the limited space between C2 and Val258 (3.4 Å). Thus this orientation of the substrate makes C1 and C6 the sites of oxidation, rather than C1 and C2. This probably explains why 4,5-dichlorocatechol is the product rather than 3,4-dichlorocatechol. Thus, structure information provides an explanation for the regiospecificity of the 3,4-DCNB dioxygenase catalysed hydroxylation.

dcnC encodes a 4,5-dichlorocatechol 1,2-dioxygenase

In the genome of strain JS3050, dcnC is the only gene similar to catechol (or chlorocatechol) 1,2-dioxygenase. To determine the function of DcnC, the expression vector pET28a-dcnC was constructed to express dcnC. The plasmid was



Fig. 3. The structure prediction of the active site of the 3,4-DCNB dioxygenase. A. Overall structure of 3,4-DCNB dioxygenase model. B, C. Model of the active site of 3,4-DCNB dioxygenase. The active site residues are shown in stick and 3,4-DCNB is in cyan. The mononuclear iron center (large orange sphere) is coordinated with two histidines and one aspartic residue. The picture was made using PyMOL (http://www.pymol.org). [Color figure can be viewed at wileyonlinelibrary.com]

Table 1. Substrate specificity of DcnC.

Substrate	Activity of cell extract of strain JS3050 (U mg^{-1})[b]	Activity of cell extract of E. coli pET28a-dcnC (U mg^{-1})[b]	Relative activity of cell extract of strain JS3050 (%)	Relative activity of cell extract of E. coli pET28a-dcnC (%)	K_m (μM)	k_{cat} (min^{-1})	k_{cat}/Km ($min^{-1} \times \mu M^{-1}$)
	1.66 ± 0.04	0.41 ± 0.01	100.0	100.0	8.0 ± 0.68	163.9	20.5
 Catechol	1.45 ± 0.04	0.42 ± 0.02	88.0	102.0	9.31 ± 0.69	144.4	15.5
 3-Methylcatechol	2.33 ± 0.11	0.66 ± 0.12	140.8	160.3	9.29 ± 1.46	310.1	33.4
 4-Methylcatechol	0.17 ± 0.01	0.05 ± 0.00	10.0	12.0	0.44 ± 0.07	17.4	39.5
 3-Chlorocatechol	1.30 ± 0.06	0.43 ± 0.02	78.5	103.8	2.44 ± 0.16	174.1	71.3
 4-Chlorocatechol	0.050 ± 0.003	0.021 ± 0.001	3.0	5.1	ND[a]	ND[a]	ND[a]
 3,4-Dichlorocatechol	0.042 ± 0.004	0.018 ± 0.000	2.5	4.4	ND[a]	ND[a]	ND[a]
 4,5-Dichlorocatechol							

[a]ND: not detected.
[b]The substrate concentration used was 500 μM.

© 2020 Society for Applied Microbiology and John Wiley & Sons Ltd,

transformed into E. coli BL21 (DE3), resulting in recombinant strain E. coli BL21 (DE3) [pET28a-dcnC]. Upon IPTG induction, the cell extract exhibited 1,2-dioxygenase activity toward various substituted catechols (Table 1). Although 4,5-dichlorocatechol is clearly the product of the initial reaction catalysed by 3,4-DCNB dioxygenase, the relative activity of DcnC against 4,5-dichlorocatechol was lower than expected. The results were similar to those seen with the 1,2,3,4-tetrachlorobenzene utilizer Pseudomonas chlororaphis RW71, where a chlorocatechol 1,2-dioxygenase TetC exhibited a lower activity toward its natural substrate 3,4,5-trichlorocatechol and 4,5-dichlorocatechol than toward other substituted catechols (Potrawfke et al., 2001).

To determine whether DcnC is the native catechol 1,2-dioxygenase in strain JS3050, the catechol 1,2-dioxygenase activity of cell extract of 3,4-DCNB-grown strain JS3050 was also measured. The substrate range was similar to that of the heterologous expressed DcnC (Table 1). Given the fact that no other catechol 1,2-dioxygenase gene (above 30% amino acid identity with DcnC) was found by bioinformatics analysis, the similar substrate ranges and the relative substrate preference of enzymes in cell extracts of strain JS3050 and E. coli BL21 (DE3) [pET28a-dcnC] provide strong evidence for the functional role of DcnC as the 4,5-dichlorocatechol 1,2-dioxygenase in the 3,4-DCNB pathway. In order to verify the affinity for different substrates to DcnC, the protein was purified (Fig. S3) and the K_m values

Theoretically, a ring-hydroxylating dioxygenase could attack 3,4-DCNB at either the 1,2- or 1,6- positions (Palatucci et al., 2019). Heterologous expression of the functional nitroarene dioxygenase from strain JS3050 established the high regiospecificity of the initial attack on 3,4-DCNB that would avoid misrouting of the resultant dichlorocatechol and ensure proper operation of the downstream pathway.

Based on the sequence information, the putative DcnR is similar to NtdR (with three amino acid differences) from Acidovorax sp. JS42 (Lessner et al., 2003) and DntR (with four amino acid differences) from Burkholderia sp. DNT (de las Heras et al., 2011). It is noteworthy that the five residue difference in NtdR (Acidovorax sp. JS42), in contrast to NagR (Ralstonia sp. strain U2), endowed NtdR with the ability to accept a number of nitroarenes as inducers. Exchange of these five residues revealed that residues at positions 227, 232 and 169, which are located near the predicted inducer-binding pocket, were key for the recognition of new compounds (Ju et al., 2009). Preliminary experiments with real-time RT-PCR indicate substantial upregulation of dcnAc by 3,4-DCNB (data not shown), which suggests inducibility of the nitroarene dioxygenase in strain JS3050 consistent with findings for 2-nitrotoluene dioxygenase NtdAcAd in JS42 (Lessner et al., 2003). This contrasts with the case in Pseudomonas stutzeri ZWLR2-1, where constitutive expression of cnbAcAd for 2-chloronitrotoluene dioxygenase was observed (Gao et al., 2016). Details of the regulation in strain JS3050 are currently under investigation.

DcnAc was aligned with α subunits of other naphthalene/nitroarene dioxygenases to compare the conserved and variable regions of the different enzymes as shown in Fig. S5. The [2Fe-2S] center binding sites are conserved in all of the enzymes. A phylogenetic tree of aromatic ring hydroxylating dioxygenases (Fig. S6) reveals that naphthalene/nitroarene dioxygenases from Gram-negative strains belong to group III aromatic ring hydroxylating dioxygenases (Parales and Resnick, 2006), which include the naphthalene dioxygenase from Pseudomonas sp. strain NCIB 98164 (Ensley et al., 1982; Suen and Gibson, 1993). The results from the amino acid sequence alignment (Fig. S6) as well as Swiss-Model structural model indicate residues in the active sites of 3,4-DCNB dioxygenase and 2,4-dinitrotoluene dioxygenase (Suen and Spain, 1993) are highly similar, particularly 3,4-DCNB dioxygenase/2,4-dinitrotoluene dioxygenase residues Gln293/Gln298, Val258/Val263, Ser350/Thr355, Ile204/ Ile209 and Ile207/Ile212. These two enzymes likely share common substrate binding orientation and substrate preference. The relationship between the amino acid residues and the substrate specify has been studied in nitrobenzene dioxygenase (Ju and Parales, 2006) and 2-nitrotoluene dioxygenase (Lee et al., 2005; Ju and Parales, 2011; Mahan et al., 2015). Future studies on DcnAc of strain JS3050 would provide additional insight about structure–function relationships.

Some of the metabolic clusters responsible for degradation of chlorinated aromatic compounds are globally dispersed and inherited through horizontal transfer of the catabolic plasmids (Wu et al., 2015). There are a number of evolutionary clues to the assembly of the functional 3,4-DCNB catabolic pathway in strain JS3050. ORFs 1, 2, 4 and 6 in the dcnDEF cluster contain genes encoding transposases and integrases found on catabolic plasmids in other isolates. They include a transposase gene similar to the one from the linuron and 3,4-dichloroaniline degrading Variovorax sp. WDL1 (Bers et al., 2013). Thus, the recruitment of dcnC might have been a key step in the assembly of the catabolic pathway for 3,4-DCNB.

The α subunit (Ac) is the most important subunit in the three component-nitroarene dioxygenase because it catalyses the dihydroxylation reaction and determines the substrate specificity (Parales et al., 1998). The large subunit DcnAc of 3,4-DCNB dioxygenase in this study is most similar (92% identity) to DntAc of 2,4-dinitrotoluene dioxygenase (Suen et al., 1996) rather than CnbAc of 2-chloronitrobenzene dioxygenase (88% identity) (Liu et al., 2011), although the latter is involved in chloronitroarene degradation (Fig. S6). Interestingly, the chemical structure of 2,4-dinitrotoluene is also similar to the 3,4-DCNB structure. Compared with the reactive nitro group, the two chlorine atoms in 3,4-DCNB are in the same position as the methyl and nitro groups in 2,4-dinitrotoluene.

The catechol and chlorocatechol 1,2-dioxygenases are historically distinguished as type I and type II enzymes, based on their substrate ranges (Dorn and Knackmuss, 1978a). Type I enzymes primarily catalyse conversion of catechol, whereas type II enzymes have a wider substrate range and transform chlorinated catechols more rapidly than catechol. However, type II chlorocatechol 1,2-dioxygenases are strongly inhibited by halocatechols simultaneously substituted in positions 4 and 5 (Haigler et al., 1988; Potrawfke et al., 2001). In the chlorobenzoate-degrading strain Pseudomonas putida AC27, 4,5-dichlorocatechol was a competitive inhibitor for the chlorocatechol 1,2-dioxygenase (Broderick and O'Halloran, 1991). DcnC, the plasmid-encoded 4,5-dichlorocatechol dioxygenase in this study and TetC (type III catechol 1,2-dioxygenase) from Pseudomonas chlororaphis RW71 (Potrawfke et al., 2001) are 95% identical, and are among the few chlorocatechol 1,2-dioxygenases reported to be active against 4,5-dichlorocatechol. Thus, DcnC in this study is a new example of type III catechol 1,2-dioxygenase.

Within the proposed chloromuconate catabolic cluster (Fig. 4B), DcnD, DcnE and DcnF exhibit high similarity (more than 60% in amino acid sequences) to the chloromuconate cycloisomerase from Variovorax sp. WDL1

© 2020 Society for Applied Microbiology and John Wiley & Sons Ltd,

Table 2. Strains and plasmids used in this study.

	Characteristics	Source or reference
Strains		
Diaphorobacter sp. strain JS3050	3,4-dichloronitrobenzene utilizer	Palatucciet al. (2019)
E. coli strains DH5α	supE44 lacU169 (ϕ80dlacZ ΔM15) recA1 endA1 hsdR17 thi-1 gyrA96 relA1	Woodcock et al. (1989)
E. coli BL21(DE3)	F^- ompT hsdS_B(r_B^- m_B^-) gal dcm lacY1 (DE3)	Novagen
Plasmids		
pUC18	Ap^r; cloning and expression vector	TaKaRa Bio
pET28a	Expression vector, Kan^r	Novagen
pUC18cnbA	pUC18 carrying cnbAaAbAcAd which encodes a 2CNB dioxygenase	Liu et al. (2011)
pUC18DcnAaAb(cnbAcAd)	Intermediate plasmid: cnbAaAb of pUC18cnbA was replaced by dcnAaAb fragment	This study
pUC18-dcnAaAbAcAd	CnbAcAd of the intermediate plasmid pUC18dcnAaAb(cnbAcAd) was replaced by DcnAcAd fragment	This study
pET28a-dcnC	PCR amplicon of dcnC cloned into pET28a	This study
pET28a-HisdcnC	PCR fragment of dcnC cloned into pET28a	This study

(Bers et al., 2013), chlorodienelactone hydrolase from Delftia acidovorans LME1 (Krol et al., 2012) and chloromaleylacetate reductase from Burkholderia sp. M701 (Sakai et al., 2014) respectively. These three enzymes DcnDEF are proposed to catalyse the sequential reactions in 3,4-dichloromuconate catabolism (Fig. 4 and Table S1), but they remain to be expressed and characterized.

pDCNB in strain JS3050 has high similarity (>90% both in query coverage and percent identity) to several previously reported catabolic plasmids including the 60.7 kb plasmid in the chlorpyrifos degrader Cupriavidus nantongensis strain X1 (Fang et al., 2019), the 77.2 kb plasmid pENH91 in the 3-chlorobenzoate degrader Cupriavidus necator strain NH9 (Ogawa and Miyashita, 1999), the 98.2 kb plasmid pA81 in the 2-chlorobenzoate degrader Achromobacter xylosoxidans A8 (Strnad et al., 2011), the 91.2 kb plasmid pCNB from the 4-chloronitrobenzene degrader Comamonas strain CNB-1 (Wu et al., 2005) and the 74.3 kb plasmid pC1-1 in the 3-chloroaniline degrader Delftia acidovorans strain CA28 (Boon et al., 2001). Notably, as shown in Fig. S7, dcnC-related genes are present in some of the plasmids (boxed). These plasmids are all larger than that of strain JS3050 but share the main plasmid backbone and some putative chlorocatechol degradation clusters with the JS3050 plasmid. This suggests all these plasmids may have a common evolutionary history. The IncP-type conjugative transfer protein Trb clusters are conserved on all these plasmids. They are broad host range and self-transmissible and could play an important role in horizontal gene transfer (Dennis, 2005).

There is an incomplete chloromuconate cycloisomerase gene that is truncated by an IS1070 adjacent to dcnC in strain JS3050. This 134-bp fragment and thetetD gene next to tetC in strain Pseudomonas chlororaphis RW71 are identical which suggests a recent common origin and that strain JS3050 may have acquired dcnC via horizontal gene transfer. During 3,4-DCNB degradation by strain JS3050, the low specific activity of DcnC toward 4,5-dichlorocatechol might be a bottleneck for the pathway and would account for the relatively slow growth on 3,4-DCNB. In addition, non-contiguous distribution of the genes responsible for 3,4-DCNB catabolism (Fig. 4B) suggests that the catabolic pathway is in the early stages of evolution.

Materials and methods

Bacterial strains, plasmids, culture conditions and chemicals

The bacterial strains and plasmids used in this study are listed in Table 2. Escherichia coli strains were grown in lysogeny broth (LB) at 37°C. Diaphorobacter sp. strain JS3050 was grown in 1/5 trypticase soy broth or mineral salts broth (Cohen-Bazire et al., 1957) medium supplied with 3,4-DCNB (1–2 mM) and Amberlite XAD-7 resin (3.5 g L^{-1}) (from Sigma-Aldrich) at 30°C. For solid media, Bacto-Agar was added to a final concentration of 12 g L^{-1}. When required, antibiotics were used at the following concentrations: ampicillin (100 mg L^{-1}), kanamycin (50 mg L^{-1}) and tetracycline hydrochloride (20 mg L^{-1}). 3,4-DCNB was purchased from Alfa Aesar (Shanghai, China); 3-chloronitrobenzene, 4-chloronitrobenzene, nitrobenzene, 2,4-dinitrotoluene, 2-nitrobenzyl alcohol, 3-nitrobenzyl alcohol, 2-nitrotoluene, 3-nitrotoluene and 4-nitrotoluene were purchased from Sinopharm Chemical Reagent (Beijing; China) with purity of higher than 98%; catechol, IPTG, 3-methylcatechol, 4,5-dichlorocatechol and 4-chlorocatechol were purchased from Sigma (St. Louis,

MO, USA); 2-chloronitrobenzene and 4-methylcatechol were from Fluka Chemical (Buchs, Switzerland); 3-chlorocatechol was from TCI (Tokyo, Japan); 3,4-dichlorocatechol was from CFW Laboratories (Walnut, CA). All the primers were synthesized by Tsingke, Shanghai, China and are listed in Table 3.

Genome sequencing and bioinformatics

DNA was extracted from cells of Diaphorobacter sp. strain JS3050 using a Mo Bio Ultra Clean microbial DNA purification kit, and sent for Illumina and PacBio sequencing by BGI Genomics (Shenzhen, China). Assembled sequence results were annotated in RAST Artemis online, and analysed for genes involved in nitroarene and chlorocatechol metabolism. Candidate genes were compared with the top biochemically characterized hits in the NCBI database using BLASTp. The sequences of the genomic DNA and pDCNB are available under accession numbers CP053553 and CP053554 respectively.

General DNA manipulations

Plasmid pUC18-dcnAaAbAcAd was constructed by replacing the cnbAaAb of pCNBA (Liu et al., 2011) with dcnAaAb and replacing the cnbAcAd with dcnAcAd. The primers used were listed in Table 3. pCNBA is a pUC18-based construct containing 2-chloronitrobenzene

constructed with Nde I digested plasmid pET28a and dcnC fragment amplified by primer pair DcnbC ndei ins/DcnbC ndei ina. The word 'in' means that the primers were used for the 'infusion clone' strategy; the characters 's' and 'a' mean 'sense' or 'antisense' of the primers respectively, 'vector' means that the primers were used to amplify the plasmid. The plasmids and inserted fragments were linked by NovoRec PCR One Step Directed Cloning Kit (Novoprotein, China) or ClonExpress II One Step Cloning Kit (Vazyme China). The molecular constructs were further confirmed by sequencing (General Biosystems, China). Other molecular operations followed standard protocols (Sambrook et al., 1989).

Biotransformation of 3,4-DCNB by heterologously expressed 3,4-DCNB dioxygenase

Plasmid pUC18-dcnAaAbAcAd was transformed into E. coli strain DH5α to construct E. coli strain DH5α [pUC18-dcnAaAbAcAd]. The cells were incubated overnight at 37°C in 5 ml of LB medium supplied with ampicillin, then transferred (1% inoculum) into 100 ml of the same medium and incubated at 30°C for 3 h. When the OD_{600} of the culture was approximately 0.6, IPTG was added (final concentration of 1 mM) the culture was incubated for another 3 h to induce the expression of dcnAaAbAcAd . Cells were collected by centrifugation, washed with sodium phosphate buffer (PB) (pH = 7.4, 50 mM) and suspended in 5 ml PB, 3,4-DCNB was added (0.15 mM final), the suspension was incubated at 30°C, and samples were removed for nitrite analysis at appropriate intervals. Escherichia coli strain DH5α [pUC18] was used as the negative control. To determine the substrate range of 3,4DCNB dioxygenase, a series of substrates (3,4-dichloronitrobenzene, nitrobenzene, 3-nitrotoluene, 2-nitrotoluene, 4-nitrotoluene, 2-chloronitrobenzene, 3-chloronitrobenzene, 4-chloronitrobenzene, 2,4-dinitrotoluene, 2,6-dinitrotoluene) were also added from 100 mM methanol solution individually for analysis. Thefinal concentrations were 1 mM. Biotransformations were performed in a shaker (180 rpm) at 30°C with cell densities of 8–10 at OD_{600}. The activities were calculated by the nitrite concentrations at 15 min for reactions against each substrate. The data are shown in Table S2.

Enzyme assays

To determine the 4,5-dichlorocatechol 1,2-dioxygenase activity of DcnC, plasmid pET28a-dcnC was transformed into E. coli BL21 (DE3) to construct strain E. coli BL21(DE3) [pET28a-dcnC]. The strain was grown in LB medium supplied with kanamycin at 30°C. When the OD_{600} reached 0.6, IPTG was added to a final concentration of 0.2 mM and the culture was incubated with shaking for another 3 h to induce the expression of dcnC. Then the cells were collected by centrifugation, washed, suspended in PB buffer and sonicated in an ice-water bath at 3.0 s on and 7.0 s off for 20 min at 30% of full power. Cell debris was removed by centrifugation at 17 000g for 60 min at 4°C. A Lambda 25 spectrophotometer (PerkinElmer/Cetus, Norwalk, CT) was used to determine the 4,5-dichlorocatechol 1,2-dioxygenase activity by a spectrophotometric assay as described previously except that EDTA was omitted (Spain and Nishino, 1987). The initial activity of DcnC against the substituted catechols was assayed by measuring the increase of OD_{260} due to the formation of corresponding muconate. The initial concentration of catechols was 500 or 50 μM. The molar extinction coefficients of muconate and its derivatives 2-methylmuconate, 3-methylmuconate, 2-chloromuconate, 3-chloromuconate, 2,3-dichloromuconate are 16 800,18 000, 13 900, 17 100, 12 400 and 12 400 M^{-1} cm^{-1} respectively (Dorn and Knackmuss, 1978b). To measure the molar extinction coefficient of 3,4-dichloromuconate, 50 μM 4,5-dichlorocatechol was incubated with excess crude cell extract until the reaction was complete to measure the increase of OD_{260}. The molar extinction coefficient of 3,4-dichloromuconate was calculated as 11 000 M^{-1} cm^{-1}. One unit of enzyme activity (U) is defined as the amount of the enzyme required for the production of 1 μmol of product in 1 min at 30°C (Wang et al., 2019). Specific activities are expressed as units per milligram of protein. Protein concentrations were measured according to the Bradford method (Bradford, 1976). For further determination of kinetic parameters the DcnC expressed by E. coli BL21(λDE3) [pET28a-HisdcnC] was purified as described previously (Zhao et al., 2018). A single colony of E. coli BL21(λDE3) [pET28a-HisdcnC] was cultured in 5 ml of fresh LB medium containing kanamycin, and was incubated in a shaker at 37°C for 14 h. Then 500 μl of bacterial culture was transferred to 500 ml of fresh LB medium and cultured at 37°C to 0.8 (OD_{600}). IPTG with a final concentration of 0.3 mM was added and cells were inculcated at 30°C for 5 h. The culture was centrifuged8000g) to collect the cells, which were resuspended with lysate buffer: pH 8.0 Tris–HCl (20 mM), 100 mM NaCl, 5% (vol./vol.) glycerol, 1 mM PMSF (phenylethanesulfonyl fluoride, protease inhibitors). The suspended cells were sonicated in an ice bath for 3.0-s on and 5.0-s off intervals for 150 cycles at 300 W and then centrifuged at 15 000g for 30 min to remove the cell debris. The supernatant was filtered through a 0.45 μM filter membrane and then loaded onto a 5-ml HisTrap HP column (GE Healthcare) and ÄKTA start system (GE Healthcare) according to the manufacturer's instructions. Two buffers were used for the protein purification. Buffer A: 20 mM Tris–HCl 8.0, 100 mM NaCl, 5% glycerol. Buffer B: 20 mM Tris–HCl 8.0, 100 mM NaCl, 500 mM imidazole, 5% glycerol. HisTrap HP column was eluted to equilibrium state with buffer A, and then 30 ml of the sample was loaded at the flow rate of 1 ml min^{-1}. Then the HisTrap HP column was

eluted with 5% and 10% buffer B. Finally, the target protein was eluted with 100% buffer B and collected. The protein purified was assessed by SDS-PAGE.

Analytical methods

GC–MS analyses were performed on a TRACE 1310 gas chromatograph (Thermo Fisher Scientific, MA, USA) using a capillary column HP-5MS (0.25 mm × 30 m, Agilent Technologies, CA, USA). The column temperature gradient was 0–2 min, 50°C; 2–11 min, 50–140°C; 11–14 min, 140°C; 14–17.5 min, 140–250°C; 17.5–20 min, 250°C. The detector was a TSQ8000 Triple Quadrupole MS (Thermo Fisher Scientific). The biotransformation samples were extracted with diethyl ether. The extracts were dried over anhydrous Na_2SO_4 before GC–MS analysis.

HPLC analysis of nitroaromatic substrates and their oxidation products was performed on a Waters e2695 Separation Module equipped with a C18 reversed-phase column (5 μm, 4.6 × 250 mm) maintained at 30°C. The monitoring wavelength was 280 nm. The mobile phase consisted of water containing 0.1% (vol./vol.) acetic acid (A) and methanol (B) at a 1.0 ml min^{-1} flow rate. The solvent gradient was 20% to 90% B over 15 min. Under these conditions, authentic 3,4-dichlorocatechol and 4,5-dichlorocatechol had retention times of 13.4 and 14.0 min respectively. The concentration of nitrite was assayed as described previously (Lessner et al., 2002), using a BioTek Synergy 2 microplate reader and sodium nitrite as a standard.

Homology modelling

The amino acid sequence of the DcnAc subunit of 3,4-DCNB dioxygenase was submitted to Swiss-model (Waterhouse et al., 2018), for homology modelling. The α subunit of 3-NT dioxygenase from Diaphorobacter sp. strain DS2 (pdb number: 5xbp), with an 87% amino acid sequence identity with the alpha subunit of 3,4-DCNB dioxygenase, was used as the modelling template (Kumari et al., 2017). The GMQE and QMEAN scores of 0.98 and −1.17 respectively showed the reliability of this model. The molecular docking process was performed with Autodock vina (Trott and Olson, 2010). The best pose with a binding affinity of −6.0 kcal mol^{-1} was used for our analysis. Based on the analysis above, PyMOL v1.3 was used to view and analyse the binding of the enzyme and 3,4-DCNB.

Acknowledgements

This work was funded by the National Natural Science Foundation of China (NSFC) (31870084, 31700093 and 31900075), DuPont Corporate Remediation Group contract LBIO-65019, and the National Key R&D Program of China (2018YFA0901200).

References

An, D.M., Gibson, D.T., and Spain, J.C. (1994) Oxidative release of nitrite from 2-nitrotoluene by a 3-component enzyme-system from Pseudomonas sp strain JS42. J Bacteriol 176: 7462–7467.

Bers, K., Batisson, I., Proost, P., Wattiez, R., De Mot, R., and Springael, D. (2013) HylA, an alternative hydrolase for initiation of catabolism of the phenylurea herbicide linuron in Variovorax sp strains. Appl Environ Microbiol 79: 5258–5263.

Boon, N., Goris, J., De Vos, P., Verstraete, W., and Top, E. M. (2001) Genetic diversity among 3-chloroaniline- and aniline-degrading strains of the Comamonadaceae. Appl Environ Microbiol 67: 1107–1115.

Booth, G. (2012) Nitro compounds, aromatic. In Ullmann's Encyclopedia of Industrial Chemistry, Weinheim: Wiley-VCH Verlag GmbH & Co. KGaA.

Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72: 248–254.

Broderick, J.B., and O'Halloran, T.V. (1991) Overproduction, purification, and characterization of chlorocatechol dioxygenase, a non-heme iron dioxygenase with broad substrate tolerance. Biochemistry 30: 7349–7358.

Cohen-Bazire, G., Sistrom, W.R., and Stanier, R.Y. (1957) Kinetic studies of pigment synthesis by non-sulfur purple bacteria. J Cell Comp Physiol 49: 25–68.

de las Heras, A., Chavarria, M., and de Lorenzo, V. (2011) Association of dnt genes of Burkholderia sp DNT with the substrate-blind regulator DntR draws the evolutionary itinerary of 2,4-dinitrotoluene biodegradation. Mol Microbiol 82: 287–299.

Dennis, J.J. (2005) The evolution of IncP catabolic plasmids. Curr Opin Biotechnol 16: 291–298.

Dorn, E., and Knackmuss, H.J. (1978a) Chemical structure and biodegradability of halogenated aromatic compounds. Two catechol 1,2-dioxygenases from a 3-chlorobenzoate-grown pseudomonad. Biochem J 174: 73–84.

Dorn, E., and Knackmuss, H.J. (1978b) Chemical structure and biodegradability of halogenated aromatic compounds. Substituent effects on 1,2-dioxygenation of catechol. Biochem J 174: 85–94.

Ensley, B.D., Gibson, D.T., and Laborde, A.L. (1982) Oxidation of naphthalene by a multicomponent enzyme system from Pseudomonas sp. strain NCIB 9816. J Bacteriol 149: 948–954.

Fang, L., Shi, T., Chen, Y., Wu, X., Zhang, C., Tang, X., et al. (2019) Kinetics and catabolic pathways of the insecticide chlorpyrifos, annotation of the degradation genes, and characterization of enzymes TcpA and Fre in Cupriavidus nantongensis X1(T). J Agric Food Chem 67: 2245–2254.

Field, J.A., and Sierra-Alvarez, R. (2008) Microbial degradation of chlorinated benzenes. Biodegradation 19: 463–480.

© 2020 Society for Applied Microbiology and John Wiley & Sons Ltd,

Fuenmayor, S.L., Wild, M., Boyes, A.L., and Williams, P.A. (1998) A gene cluster encoding steps in conversion of naphthalene to gentisate in Pseudomonas sp. strain U2. J Bacteriol 180: 2522–2530.

Gao, Y.Z., Liu, H., Chao, H.J., and Zhou, N.Y. (2016) Constitutive expression of a nag-like dioxygenase gene through an internal promoter in the 2-chloronitrobenzene catabolism gene cluster of Pseudomonas stutzeri ZWLR2-1. Appl Environ Microbiol 82: 3461–3470.

Gao, Y.Z., Liu, X.Y., Liu, H., Guo, Y., and Zhou, N.Y. (2020) A Bph-like nitroarene dioxygenase catalyzes the conversion of 3-nitrotoluene to 3-methylcatechol by Rhodococcus sp. strain ZWL3NT. Appl Environ Microbiol 86(4), e02517–19.

Haigler, B.E., Nishino, S.F., and Spain, J.C. (1988) Degradation of 1,2-dichlorobenzene by a Pseudomonas sp. Appl Environ Microbiol 54: 294–301.

Johnson, G.R., Jain, R.K., and Spain, J.C. (2002) Origins of the 2,4-dinitrotoluene pathway. J Bacteriol 184: 4219–4232.

Ju, K.S., Parales, J.V., and Parales, R.E. (2009) Reconstructing the evolutionary history of nitrotoluene detection in the transcriptional regulator NtdR. Mol Microbiol 74: 826–843.

Ju, K.S., and Parales, R.E. (2006) Control of substrate specificity by active-site residues in nitrobenzene dioxygenase. Appl Environ Microbiol 72: 1817–1824.

Ju, K.S., and Parales, R.E. (2009) Application of nitroarene dioxygenases in the design of novel strains that degrade chloronitrobenzenes. Microb Biotechnol 2: 241–252.

Ju, K.S., and Parales, R.E. (2010) Nitroaromatic compounds, from synthesis to biodegradation. Microbiol Mol Biol Rev 74: 250–272.

Ju, K.S., and Parales, R.E. (2011) Evolution of a new bacterial pathway for 4-nitrotoluene degradation. Mol Microbiol 82: 355–364.

Katsivela, E., Wray, V., Pieper, D.H., and Wittich, R.M. (1999) Initial reactions in the biodegradation of 1-chloro-4-nitrobenzene by a newly isolated bacterium. Appl Environ Microbiol 65: 1405–1412.

Krol, J.E., Penrod, J.T., McCaslin, H., Rogers, L.M., Yano, H., Stancik, A.D., et al. (2012) Role of IncP-1beta plasmids pWDL7::rfp and pNB8c in chloroaniline catabolism as determined by genomic and functional analyses. Appl Environ Microbiol 78: 828–838.

Kumari, A., Singh, D., Ramaswamy, S., and Ramanathan, G. (2017) Structural and functional studies of ferredoxin and oxygenase components of 3-nitrotoluene dioxygenase from Diaphorobacter sp strain DS2. PLoS One 12: 16.

Lee, K.S., Parales, J.V., Friemann, R., and Parales, R.E. (2005) Active site residues controlling substrate specficity in 2-nitrotoluene dioxygenase from Acidovorax sp. strain JS42. J Ind Microbiol Biotechnol 32: 465–473.

Lessner, D.J., Johnson, G.R., Parales, R.E., Spain, J.C., and Gibson, D.T. (2002) Molecular characterization and substrate specificity of nitrobenzene dioxygenase from Comamonas sp strain JS765. Appl Environ Microbiol 68: 634–641.

Lessner, D.J., Parales, R.E., Narayan, S., and Gibson, D.T. (2003) Expression of the nitroarene dioxygenase genes in Comamonas sp strain JS765 and Acidovorax sp strain JS42 is induced by multiple aromatic compounds. J Bacteriol 185: 3895–3904.

Liu, H., Wang, S.J., Zhang, J.J., Dai, H., Tang, H.R., and Zhou, N.Y. (2011) Patchwork assembly of nag-like nitroarene dioxygenase genes and the 3-chlorocatechol degradation cluster for evolution of the 2-chloronitrobenzene catabolism pathway in Pseudomonas stutzeri ZWLR2-1. Appl Environ Microbiol 77: 4547–4552.

Mahan, K.M., Penrod, J.T., Ju, K.S., Al Kass, N., Tan, W.A., Truong, R., et al. (2015) Selection for growth on 3-nitrotoluene by 2-nitrotoluene-utilizing Acidovorax sp strain JS42 identifies nitroarene dioxygenases with altered specificities. Appl Environ Microbiol 81: 309–319.

Nishino, S.F., and Spain, J.C. (1995) Oxidative pathway for the biodegradation of nitrobenzene by Comamonas sp strain JS765. Appl Environ Microbiol 61: 2308–2313.

Ogawa, N., and Miyashita, K. (1999) The chlorocatechol-catabolic transposon Tn5707 of Alcaligenes eutrophus NH9, carrying a gene cluster highly homologous to that in the 1,2,4-trichlorobenzene-degrading bacterium Pseudomonas sp. strain P51, confers the ability to grow on 3-chlorobenzoate. Appl Environ Microbiol 65: 724–731.

Palatucci, M.L., Waidner, L.A., Mack, E.E., and Spain, J.C. (2019) Aerobic biodegradation of 2,3- and 3,4-dichloronitrobenzene. J Hazard Mater 378: 120717.

Parales, J.V., Parales, R.E., Resnick, S.M., and Gibson, D. T. (1998) Enzyme specificity of 2-nitrotoluene 2,3-dioxygenase from Pseudomonas sp. strain JS42 is determined by the C-terminal region of the alpha subunit of the oxygenase component. J Bacteriol 180: 1194–1199.

Parales, R.E., and Resnick, S.M. (2006) Aromatic ring hydroxylating dioxygenases. In Pseudomonas, Vol 4: Molecular Biology of Emerging Issues, pp. 287–340). New York: Springer.

Potrawfke, T., Armengaud, J., and Wittich, R.M. (2001) Chlorocatechols substituted at positions 4 and 5 are substrates of the broad-spectrum chlorocatechol 1,2-dioxygenase of Pseudomonas chlororaphis RW71. J Bacteriol 183: 997–1011.

Potrawfke, T., Timmis, K.N., and Wittich, R.M. (1998) Degradation of 1,2,3,4-tetrachlorobenzene by Pseudomonas chlororaphis RW71. Appl Environ Microbiol 64: 3798–3806.

Sakai, Y., Ogawa, N., Shimomura, TD [(Pseudo [(.6(aawfke,)-FJiuji2(Y.,)37

Strnad, H., Ridl, J., Paces, J., Kolar, M., Vlcek, C., and Paces, V. (2011) Complete genome sequence of the haloaromatic acid-degrading bacterium Achromobacter xylosoxidans A8. J Bacteriol 193: 791–792.

Suen, W.C., and Gibson, D.T. (1993) Isolation and preliminary characterization of the subunits of the terminal component of naphthalene dioxygenase from Pseudomonas putida NCIB 9816-4. J Bacteriol 175: 5877–5881.

Suen, W.C., Haigler, B.E., and Spain, J.C. (1996) 2,4-Dinitrotoluene dioxygenase from Burkholderia sp. strain DNT: similarity to naphthalene dioxygenase. J Bacteriol 178: 4926–4934.

Suen, W.C., and Spain, J.C. (1993) Cloning and characterization of Pseudomonas sp. strain DNT genes for 2,4-dinitrotoluene degradation. J Bacteriol 175: 1831–1837.

Trott, O., and Olson, A.J. (2010) AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. J Comput Chem 31: 455–461.

van der Meer, J.R., Eggen, R.I., Zehnder, A.J., and de Vos, W.M. (1991) Sequence analysis of the Pseudomonas sp. strain P51 tcb gene cluster, which encodes metabolism of chlorinated catechols: evidence for specialization of catechol 1,2-dioxygenases for chlorinated substrates. J Bacteriol 173: 2425–2434.

Wackett, L.P. (2004) Evolution of enzymes for the metabolism of new chemical inputs into the environment. J Biol Chem 279: 41259–41262.

Wang, L., Gao, Y.-Z., Zhao, H., Xu, Y., and Zhou, N.-Y. (2019) Biodegradation of 2-bromonitrobenzene by Pseudomonas stutzeri ZWLR2-1. Int Biodeter Biodegr 138: 87–h.

Waterhouse, A., Rempfer, C., Heer, F.T., Studer, G., Tauriello, G., Bordoli, L., et al. (2018) SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic Acids Res 46: W296–W303.

Woodcock, D.M., Crowther, P.J., Doherty, J., Jefferson, S., DeCruz, E., Noyer-Weidner, M., et al. (1989) Quantitative evaluation of Escherichia coli host strains for tolerance to cytosine methylation in plasmid and phage recombinants. Nucleic Acids Res 17: 3469–3478.

Wu, J.F., Jiang, C.Y., Wang, B.J., Ma, Y.F., Liu, Z.P., and Liu, S.J. (2006) Novel partial reductive pathway for 4-chloronitrobenzene and nitrobenzene degradation in Comamonas sp strain CNB-1. Appl Environ Microbiol 72: 1759–1765.

Wu, J.F., Sun, C.W., Jiang, C.Y., Liu, Z.P., and Liu, S.J. (2005) A novel 2-aminophenol 1,6-dioxygenase involved in the degradation of p-chloronitrobenzene by Comamonas strain CNB-1: purification, properties, genetic cloning and expression in Escherichia coli. Arch Microbiol 183: 1–8.

Wu, Y., Arumugam, K., Tay, M.Q., Seshan, H., Mohanty, A., and Cao, B. (2015) Comparative genome analysis reveals genetic adaptation to versatile environmental conditions and importance of biofilm lifestyle in Comamonas testosteroni. Appl Microbiol Biotechnol 99: 3519–3532.

Zeiger, E., Anderson, B., Haworth, S., Lawlor, T., Mortelmans, K., and Speck, W. (1987) Salmonella mutagenicity tests: III. Results from the testing of 255 chemicals. Environ Mol Mutagen 9: 61–109.

Zhao, H., Xu, Y., Lin, S., Spain, J.C., and Zhou, N.Y. (2018) The molecular basis for the intramolecular migration (NIH shift) of the carboxyl group during para-hydroxybenzoate catabolism. Mol Microbiol 110: 411–424.

Zhen, D., Liu, H., Wang, S.J., Zhang, J.J., Zhao, F., and Zhou, N.Y. (2006) Plasmid-mediated degradation of 4-chloronitrobenzene by newly isolated Pseudomonas putida strain ZWL73. Appl Microbiol Biotechnol 72: 797–803.

Zhou, N.Y., Al-Dulayymi, J., Baird, M.S., and Williams, P.A. (2002) Salicylate 5-hydroxylase from Ralstonia sp strain U2: a monooxygenase with close relationships to and shared electron transport proteins with naphthalene dioxygenase. J Bacteriol 184: 1547–1555.

Zhou, N.Y., Fuenmayor, S.L., and Williams, P.A. (2001) nag genes of Ralstonia (formerly Pseudomonas) sp. strain U2 encoding enzymes for gentisate catabolism. J Bacteriol 183: 700–708.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1 : supporting Information

© 2020 Society for Applied Microbiology and John Wiley & Sons Ltd,